



A Risk Analysis of *Brucella abortus* Transmission Among Bison, Elk, and Cattle in the Northern Greater Yellowstone Area



Technical Report to the National Park Service
October 2010

YCR-2012-02



Suggested citation: Yellowstone Center for Resources. 2012. A Risk Analysis of *Brucella abortus* Transmission Among Bison, Elk, and Cattle in the Northern Greater Yellowstone Area. Technical Report for the National Park Service, October 2010. Mammoth Hot Springs, Wyoming, YCR-2012-02.

A Risk Analysis of *Brucella abortus* Transmission among Bison, Elk, and Cattle in the Northern Greater Yellowstone Area

Brant A. Schumaker,^{1*} Jonna A.K. Mazet,^{2,3} John Treanor,⁴ Rick Wallen,⁴ Ian A. Gardner,³ Martin Zaluski,⁵ and Tim E. Carpenter^{1,3}

¹Center for Animal Disease Modeling and Surveillance (CADMS), University of California, Davis, California

²Wildlife Health Center, University of California, Davis, California

³Department of Medicine and Epidemiology, University of California, Davis, California

⁴National Park Service, Yellowstone National Park, Wyoming

⁵Montana Department of Livestock, Helena, Montana

October 2010

CONTENTS

Title Page	1
Table of Contents	2
Background	3
Objective 1 – Bison or Elk: Who should be the Target of Brucellosis Control in the Northern Greater Yellowstone Area?	39
Objective 2 – Who infects whom? Interspecies Transmission Dynamics of Brucellosis in the Northern Greater Yellowstone Area	73
Objective 3 – Brucellosis Management Strategies in the Northern Greater Yellowstone Area	96
Conclusions	122

BACKGROUND

Detection and Transmission Dynamics of *Brucella abortus* in the Greater Yellowstone Area

Disease management at the wildlife-livestock interface is hampered by the challenge of balancing wildlife conservation with the livelihoods and traditions of livestock producers. The potential for disease transmission between wildlife and livestock exacerbates conflicts between natural resource managers and cattlemen, reduces tolerance for wildlife near livestock operations, and negatively impacts conservation. Therefore, diseases that affect both wildlife and livestock are important in resource management, regardless of their direct impact to the wild animal populations which may serve as their reservoirs. Many important diseases of livestock are shared among multiple species, including foot-and-mouth disease, Rift Valley fever, and Johne's disease (Daszak et al., 2000; Chivian, 2001; Taylor et al., 2001; Woolhouse et al., 2001; Belloy et al., 2004; Cunningham, 2005; Böhm et al., 2009; Tomley and Shirley, 2009). Human population growth and associated landscape changes, as well as competition for grazing lands, have made wildlife-livestock disease transmission more likely by reducing the spatial separation between livestock operations and wildlife habitat (Daszak et al., 2001; Western, 2001).

The US is free of many of the devastating diseases affecting both wildlife and livestock worldwide. However, the government has spent billions of dollars on disease eradication programs for both wildlife and livestock. Pneumonia caused by multiple pathogens from domestic sheep threaten bighorn sheep populations throughout the western US (Clifford et al., 2009; USDA Forest Service, 2010). Tuberculosis in Michigan deer and cattle populations continues to be a problem (State of Michigan, 2008). Also, recent cases of tuberculosis in captive elk in Nebraska and cattle in California are an indication that the US is far from eradication of

these diseases (Olmstead and Rhode, 2004). In addition, multiple recurrences of bovine brucellosis, caused by the bacterium *Brucella abortus* in the states surrounding the greater Yellowstone area have greatly complicated the US eradication effort.

B. abortus is a gram-negative, facultative, intracellular bacterium that causes disease in many domestic and wild animal species including cattle, bison (*Bison bison*), elk (*Cervus elaphus*), and moose (*Alces alces*) (Creech, 1930; Thorne et al., 1978a; Edmonds et al., 1999). Bacteria invade the mucous membranes of ungulates and can cause placentitis with late-gestation abortions in females and orchitis and epididymitis in males (Bercovich, 1998). Increased abortion rates, decreased milk production, loss of condition, infertility, and lameness in cattle have made brucellosis extremely important to beef and milk producers around the world (Manthei and Carter, 1950), restricting international trade in many instances (Wilson and Beers, 2001).

B. abortus was first characterized as the cause of epizootic abortion by Bernard Bang in 1896 (Bang, 1897). The eradication of the disease from the US has been a priority of the federal government since 1934, when a Cooperative State-Federal Brucellosis Eradication Program (BEP) was adopted to reduce the prevalence of brucellosis in cattle, designating it as the most significant livestock disease at that time. Since then, agencies have implemented a variety of livestock, wildlife, and disease risk management strategies (Cheville et al., 1998). Billions of dollars have been spent eradicating brucellosis from livestock in nearly every state in the US (Wise, 1980).

Zoonotic Implications of Brucellosis

The potential for human infection and large economic losses have made *B. abortus* an important pathogen restricting international trade (Wilson and Beers, 2001). The bacterium has also been classified as an overlap human/livestock select agent by the United States Department of Health and Human Services and the United States Department of Agriculture (USDA) (2002). Brucellosis in humans is characterized by intermittent bacteremia caused by seeding of bacteria from lymph nodes which causes malaise, aching joints, and irregular spikes in body temperature referred to as undulant fever. The recommended treatment of human brucellosis is doxycycline and rifampin (Centers for Disease Control and Prevention, 2007).

In 2006, human brucellosis from exposure to *B. abortus* in cattle, *B. melitensis* in sheep and goats, and *B. suis* in swine was still considered the most common zoonotic infection worldwide (Pappas et al., 2006). In the early years of the BEP, human brucellosis in the US was mainly acquired from contact with infected meat and tissues during slaughter operations. However, *Brucella spp.* also colonize the mammary glands of infected animals and can be transmitted in milk (Young and Suvannoparrat, 1975). In the last two decades, the main cause of human brucellosis in the US was from food-borne infection mainly through importation of soft cheeses from Mexico (Chomel et al., 1994).

Although the US likely has low exposure to *B. abortus* in humans, the increasing number of *B. abortus* infections in Kyrgyzstan is an example of what can happen without appropriate disease control strategies. After the collapse of the Soviet Union, the country was ill-equipped to handle a major livestock disease like brucellosis. These circumstances, combined with the impacts of a depressed economy and poor hygiene, has created the opportunity for a re-emerging zoonotic disease epidemic. In the first six months of 2003, there were 1170 reported cases of

human brucellosis in Kyrgyzstan, a 30% increase from the previous year (UN Office for the Coordination of Humanitarian Affairs, 2003). Of those cases, 20% were children or adolescents. Because brucellosis is so difficult and expensive to treat it has been a great detriment to the economy of the country and gives credence to the expenditures on the BEP in the US (Kozukeev et al., 2006).

State-Federal Cooperative Brucellosis Eradication Plan

During the 76-year history of the BEP, it has limited the impacts of brucellosis in cattle throughout the US (Donch and Gertonson, 2008). At the program's inception, 11.5% of adult cattle in the US were infected with the bacterium (Ragan, 2002), and the annual losses to the livestock industry were \$400 million with \$50 million lost to decreases in milk production alone (Knox, 1947). In 1957, there were an estimated 124,000 herds infected with brucellosis using imperfect surveillance with only 33-50% detection (Ragan, 2002). By 1961, the entire annual loss to the livestock industry was reduced to \$25 million (Mingle, 1961), and by 2000 only 6 herds were diagnosed as infected. Studies have shown that, if the program were discontinued, costs would increase by \$80 million annually in less than 10 years (Bittner, 2004). By early 2008, the US and all associated territories were brucellosis-free in livestock. However in June 2008, brucellosis was again detected in cattle herds in Montana and Wyoming (Donch and Gertonson, 2008). Transmission incidents in the last four years in all three states surrounding Yellowstone National Park (YNP) – Idaho, Montana, and Wyoming – have highlighted the importance of wildlife brucellosis on livelihoods and management (Donch and Gertonson, 2008).

Greater Yellowstone Area and Wildlife Populations

YNP was established as America's first national park in 1872, and has become a flagship for wildlife conservation worldwide. Despite its large size of 8,987 square kilometers, YNP is not independent of its surrounding ecosystem, the greater Yellowstone area (GYA). The GYA is one of the largest intact temperate zone ecosystems on earth and includes approximately 28,000 square miles in Montana, Idaho, and Wyoming and encompasses state lands, two national parks, portions of six national forests, three national wildlife refuges, Bureau of Land Management holdings, and private and tribal lands. The GYA is also home to the largest wild and free-ranging elk and bison populations in the US.

Approximately 125,000 elk occupy the GYA across 25 elk management jurisdictions. Agencies manage elk and their habitat resources through complex interagency cooperation. Elk hunting occurs in all involved elk management jurisdictions except YNP. There are also 23 elk feedgrounds in northwest Wyoming (the National Elk Refuge and 22 state operations) that may support approximately 25,000 elk, depending on winter severity. Approximately 5,000 bison occupy the GYA across trans-boundary bison management jurisdictions in and adjacent to YNP (4,200) and Jackson Hole, Wyoming (800). Bison hunting presently occurs only in select national forest areas in Wyoming, with most bison in Jackson Hole utilizing the feedground on the National Elk Refuge during the winter.

At the turn of the century, only 50,000 elk were reportedly remaining in the entire continental US, mainly inhabiting areas of the GYA (Seton, 1927). Supplemental winter elk feeding began in Jackson Hole, Wyoming in 1910 as an effort to help elk avoid starvation during harsh winters and decrease their impacts on agricultural lands (Smith, 2001). This practice was expanded in 1912, by the creation of the National Elk Refuge, but supplemental feeding has

ultimately led to a number of negative consequences. Elk are above management targets in many areas in the GYA (Dickson, 2005). Feeding practices have artificially increased their population density from November to April and allowed for more intraspecies transmission of diseases during the winter months. For example, longer feeding seasons are associated with higher *B. abortus* seroprevalence (Cross et al., 2007).

However, the economics of elk hunting in Wyoming have made the possibility of closing feedgrounds extremely controversial. Unguided hunting on public lands in Wyoming is prohibited for non-residents, and the outfitting industry is a large part of the local economy. In 1980, outfitting businesses in Teton County, Wyoming had direct sales of \$2.4 million for big-game hunting (Taylor et al., 1981). Accounting for indirect revenue yielded a total of \$4.2 million in annual income from outfitting (Taylor et al., 1981). Because most hunting revenues are generated in the fall, the outfitting industry helps to bridge the gap between summer and winter tourist seasons.

The continental divide runs from west to east across the southern portion of YNP. The northern GYA includes the Yellowstone bison population and five elk populations (Gallatin-Madison, Gravelly-Snowcrest, Madison-Firehole, northern Yellowstone, and Sand Creek, Idaho), which are distributed across over 1,100 square miles in the northern GYA. Estimates of northern Yellowstone elk were near 25,000 animals in the late 1980s but decreased by approximately 50-60% by 2006 (Eberhardt et al., 2007). The Yellowstone bison population ranges between 2000 and 5000 individuals (Meagher, 1973; Clarke et al., 2005) depending on environmental conditions and management strategies implemented. These bison are important for the conservation of the species because the population is derived from the original wild herd supplemented by an introduced herd containing diverse genetics (Meagher, 1973). In addition,

the bison have had no evidence of cattle-hybridization (Halbert et al., 2005). Therefore, disease management activities, including the future potential for movement of individual bison into other herds, are of special interest in this population.

The 2009 summer count for the Yellowstone bison herd was 3,300 animals, divided equally between a central and northern breeding population. The modeled food-limiting carrying capacity for bison within YNP is 6200 individuals (Plumb et al., 2009). However, even at lower population numbers, interactive effects of severe winters and herd density with population numbers greater than 4200 have lead to large-scale dispersal to lower elevations. Plumb et al. (2009), recommended the Yellowstone bison herd be maintained with less than 4500 animals to abate most large-scale movements outside the park during average winter conditions. Appropriate population management would help avoid contact with cattle, which are grazed (266 in the winter and 1363 in the spring) on public and private lands adjacent to YNP and within habitat occupied by bison and elk during the winter (Kilpatrick et al., 2009).

Brucellosis Pathogenesis in Wildlife

The proximity of cattle-grazing to wildlife populations makes interspecies disease transmission a concern. Wild, free-ranging bison and elk in the GYA persist as the last known reservoir of *B. abortus*-caused brucellosis in the US (Godfroid, 2002). Brucellosis in Yellowstone bison is similar to that of chronically infected cattle (Roffe et al., 1999; Rhyan et al., 2001). In the wildlife host, *B. abortus* is typically transmitted to susceptible individuals after licking a newborn calf of an infected dam or ingesting an aborted fetus or placenta. Once inside the host, *B. abortus* resides in regional lymph nodes and then is transported to other lymph nodes. The sublumbar or supramammary lymph nodes are common targets. *Brucella abortus* uses

several strategies to evade detection by the host's immune system (Arenas et al., 2000). The bacterium often takes up residence in host macrophages, where its intracellular signals prevent phagosome-lysosome fusion (Frenchick et al., 1985; Pizarro-Cerda et al., 1998). During the third trimester of pregnancy, the bacterium preferentially invades the placenta and causes fetal death and abortion. Nearly 100% of bison will abort their first calf after infection (Davis et al., 1990; Davis et al., 1991). The typical clinical signs seen in the aborted material are necrotizing placentitis and fetal pneumonia (Rhyan et al., 2001). If the fetus is carried to term, the bacterium may also be vertically transmitted to the calf from the dam's milk. However, there has been no proven relationship between the serostatus of dam and calf, and most calves are seronegative by six months of age (Fuller et al., 2007).

The incubation period for brucellosis is variable and affected by gestation, exposure dose, age, vaccination, and effects of host-resistance (Nicoletti, 1980). After experimental inoculation of elk, mean length of time between inoculation and a serologic titer was 39 days, and the mean time-to-abortion was 89 days post-infection (Thorne et al., 1979). In cattle, about 20% of calves born to infected dams are seronegative but latently infected (Plommet et al., 1973; Lapraik et al., 1975). Up to 10% of these calves have been known to seroconvert in early adulthood as the stress of pregnancy lowers their immune systems (Wilesmith, 1978). This so-called "heifer syndrome" has been described in bison and elk in addition to cattle (Van Den Born and Vervoorn, 1965; Plommet et al., 1973; Lapraik et al., 1975; Thorne et al., 1978b; Thorne et al., 1979; Catlin and Sheehan, 1986; Olsen et al., 2003). Once an animal is infected, there is little evidence to suggest that it will ever recover from infection, and it is recommended that it be considered a carrier for life, even if no abortions occur (Ragan, 2002).

Brucellosis in the GYA

Brucellosis was first detected among wildlife in the GYA in 1917, when epizootic abortion was described in Yellowstone bison (Mohler, 1917). The now-bison disease was most likely acquired from domestic cattle which were brought into the area for grazing (Meagher and Meyer, 1994). Elk in the southern GYA probably acquired the disease directly from cattle and then transmitted it to the bison presently using Grand Teton National Park. Today, elk populations in the northern GYA have low seroprevalence (i.e., exposure; <5%) for *B. abortus*, whereas seroprevalence in Yellowstone bison is high (40-60%) (Cheville et al., 1998). Elk feedgrounds in the southern GYA have increased the prevalence of brucellosis. The average seroprevalence for *B. abortus* among fed elk is around 26% (Aune, 2002; Etter and Drew, 2006). Brucellosis in wildlife does not generally threaten population persistence. Coinfection with bovine tuberculosis reduced the pregnancy rates by 10-15% for bison in Wood Buffalo National Park in the absence of elk competing for grazing land (Joly and Messier, 2005). However in YNP bison, recruitment and population numbers have remained sustainable, aside from boundary removals from the population to reduce transmission risk to cattle.

Although the wildlife populations in the GYA are stable, the ability of bison and elk to concomitantly serve as alternative hosts and sources of *B. abortus* increases the complexity of risk of transmission to cattle. This multi-reservoir system poses significant challenges to comprehensive disease management (Delahay et al., 2009). Understanding the interspecies transmission dynamics of a multi-host system is crucial for disease management (Dobson, 2004; Delahay et al., 2009). Some hosts may be persistent reservoirs of disease, and others may be recurrently infected through pathogen spillover (Power and Mitchell, 2004). Overall, diseases

with multiple wildlife hosts are deemed extremely difficult to control and eradicate (Government Accountability Office, 2009).

Brucellosis Diagnostics

In order for a disease management program to be effective, infected animals must be detected. Unfortunately, there is no perfect brucellosis reference test. Although culture of tissues or fluids, such as milk, is frequently used as a standard for *B. abortus* diagnosis, it is also imperfectly sensitive (Gall and Nielsen, 2004). There are often few detectable bacteria and no obvious signs of infection (i.e., subclinical or latent infection). If an individual clears the infection, it is likely to test positive on serologic tests, yet not shed bacteria. Also, collection, handling, and storage of samples, as well as laboratory techniques can affect the success of culture (Sutherland, 1980; Rhyan et al., 1997; Roffe et al., 1999; Gall and Nielsen, 2004). Laboratory methods require specific media and specialized incubation conditions, and *B. abortus*'s slow growth rate often leads to overgrowth of non-target bacteria on the culture plates. Because of these difficulties, serologic testing is frequently used to determine infection status.

The ideal serologic test should correctly classify an animal's infection status, be able to be performed animal-side, and yield rapid results. However, under field conditions, where an individual has the opportunity for exposure to *B. abortus* organisms, it is impossible to determine whether serologic test-positive but culture-negative individuals are either exposed but not currently infected or truly infected with undetected bacteria due to the lack of sensitivity of culture. Conversely, recently infected individuals may not yet be producing sufficient antibodies for serologic detection. However, these false-negative reactors may shed bacteria when aborting or calving. Also, persistently infected individuals can falsely test negative after the immune

response diminishes below the threshold of detection due to lack of repeated exposure. Thus, it is unlikely that all truly infected individuals can be identified by serology alone (Cheville et al., 1998). Roffe et al. (1999) noted that only approximately 46% of sero-positive bison were culture positive from one or more tissues.

As the prevalence of brucellosis decreases in the US, the need for diagnostic tests with high sensitivity and specificity is becoming much more critical for appropriate brucellosis management, with the cost of incorrect test results becoming more substantial. Many serologic tests have been produced to aid diagnosis of *B. abortus* infections; however, all currently-used diagnostic methods were developed and validated for use only in cattle. When applied to wildlife, many cattle tests have been shown to be inaccurate and unpredictable (Morton et al., 1981; Davis et al., 1990). Furthermore, antibodies developed to environmental bacteria, such as *Yersinia enterocolitica* O:9, can cause cross-reactions in commonly-used *B. abortus* screening tests (Kittelberger et al., 1995).

In the last decade, some tests have shown promise in tackling these diagnostic issues. In 1999, Edmonds and colleagues described a western immunoblot designed to differentiate antibody responses to *B. abortus*, *B. melitensis*, and *B. suis*, as well as *Yersinia enterocolitica* O:9. The variation in O-antigens among the different bacterial species results in the host's development of specific antibodies to *B. abortus* that can be differentiated by the western blot (Edmonds et al., 1999). The technique was evaluated for use in detecting *B. abortus* antibodies in elk, and the results were comparable to standard serologic tests (Schumaker et al., 2010). In 2000, Gall et al. validated a fluorescence polarization assay (FPA) for use in detecting serum antibodies for *B. abortus* in bison (Gall et al., 2000). The authors estimated the specificity of FPA and other serologic tests in a population with no epidemiologic evidence of the presence of

brucellosis, and a blinded study yielded sensitivity and specificity values of 96.3% and 97.6%, respectively.

Interspecies Disease Transmission

Both elk and bison have been shown to be competent reservoirs for *B. abortus* transmission to cattle (Thorne et al., 1979; Davis et al., 1990). In an elk-cattle pen study by Thorne et al. (1978a), close confinement may have contributed to transmission, but contact was not closer than feedground situations. Also, a report by Flagg (1983) showed evidence of fence-line contact and transmission of *B. abortus* from bison to cattle. While *B. abortus* may be carried in sperm and transmission via artificial insemination is a concern in livestock, males are not considered to be an important source of transmission risk from wildlife to cattle (Thorne, 2001).

The risk period for *B. abortus* transmission is well-defined. In general, data suggest that bison and elk in the northern portion of the GYA exhibit a high degree of birth synchrony, with the majority (80%) of bison calving during late-April to late-May and elk calving between mid-May to mid-June (Cheville et al., 1998; Berger and Cain, 1999). Feed ground data from the southern portion of the GYA in Wyoming have shown birth dates for elk that are later in the year, but parturition events are still unlikely after the third week of June due to the normal pattern of sexual segregation (Cross et al., 2009; Maichak et al., 2009). Including abortions in the last 90 days of pregnancy, late-January to mid-June is the most likely period for *B. abortus* transmission (Roffe et al., 2004).

The probability of *B. abortus* transmission between elk (or from elk to cattle) is likely low during calving (May through June) because elk dams segregate themselves while giving birth and meticulously clean the birth site (Johnson, 1951). Thus, birth sites are dispersed, and

the likelihood of other elk or cattle encountering infected birth tissues and fluids is low. However, transmission risk may be higher during the brucellosis abortion period from February through April when many elk aggregate in larger groups on lower-elevation winter ranges that sometimes include ranch areas with cattle (Hamlin and Cunningham, 2008). Spontaneous abortions by elk that are not segregated from the herd could expose many susceptible elk (or cattle) to infected fetuses and birth tissues (P.J. White, personal communication). Elk that winter at the Madison headwaters showed 53% winter range overlap with Yellowstone bison in December and 76% overlap in May (Ferrari and Garrot, 2002). A meaningful percentage of elk locations (18%) were within 100 meters of bison with comingling correlated with snowpack. However, these elk do not show evidence of an increase in *B. abortus* exposure compared with populations with spatio-temporal separation from bison (Ferrari and Garrot, 2002; Proffitt et al., 2010).

In contrast to elk, bison are gregarious during parturition, and pregnant females have been observed to nuzzle newborn calves (Treanor et al., 2008). Mobbing events of a newborn calf or aborted fetus could contribute to intra-species transmission of *B. abortus* if the dam were infected (Jones et al., 2009). Bison conservation continues to be a priority of the National Park Service; however, for decades, livestock and regulatory personnel have viewed Yellowstone bison as a potential source of pathogens for livestock in the GYA (Meagher and Meyer, 1994). Current management, which maintains spatial and temporal separation between bison and cattle, makes the risk of *B. abortus* transmission from bison to cattle in the northern GYA negligible (Kilpatrick et al., 2009). However, hazing and culling actions by bison managers to maintain this separation have been highly scrutinized and criticized for their economic costs and negative effects to bison. In the last decade, there have been multiple detections of brucellosis in cattle in

the GYA states (Idaho, Montana, Wyoming), with elk identified as the source of infection for nine cases since 2002 (Donch and Gertonson, 2008).

It is unknown how close a susceptible cow would have to be to *B. abortus*-infected tissues before it would be likely to investigate tissues and become exposed to the bacterium. The Starkey Project at the Pacific Northwest Research Station found that forage competition between elk and cattle likely decreases the chance of comingling on winter range (Coe et al., 2005). However, lack of available forage and other environmental pressures during severe winters in the GYA might increase comingling. The number of days a *B. abortus*-contaminated birth site is infective is dependent upon the amount of time that it takes for an infected fetus or tissues to be scavenged or for ultraviolet radiation to degrade the bacteria. Aune et al. (2007) and Cook et al. (2004) found that fetuses would be scavenged prior to ultraviolet degradation of bacteria (mean; range = 18.2; 1-78 days), which was used by Kilpatrick et al. (2009) to estimate the persistence of an infected site.

Brucellosis Status in GYA States

In the last four years, there have been outbreaks of brucellosis in all three states in the GYA (Idaho, Montana, and Wyoming). Idaho lost its brucellosis class-free status in 2006 but regained it in 2007. It has been maintaining a surveillance boundary for the five counties bordering YNP (USAHA, 2009). Idaho requires mandatory official calfhood vaccination for all cattle operations. It has been working to develop herd plans to minimize the risk of *B. abortus* transmission from wildlife to cattle by mainly focusing on preventing winter feeding of elk, fencing stack yards, securing hay barns, and enclosing winter cattle feedgrounds.

Montana had cattle herds test positive for brucellosis in May, 2007 and June, 2008, causing the state to lose its brucellosis-free status. The state regained its class-free status in July, 2009 and continued its surveillance and Brucellosis Action Plan (BAP) until January 10th, 2010 (6 months following reclassification to class-free status). The high-risk area for the BAP, included the seven counties surrounding YNP (Beaverhead, Madison, Park, Gallatin, Sweet Grass, Stillwater, Carbon). After January, 2010 the more recent risk area included only Beaverhead, Madison, Park, and Gallatin counties. Surveillance of elk provided 880 useable samples, of which 62 (7%) were positive on standard serologic tests. However, only 13 (1.5% of the total) were confirmed positive by the western blot. Seropositive elk were found in five distinct hunting districts.

Wyoming lost its brucellosis-free status in 2004 and regained it in 2006. It had one herd test positive in 2008 but the outbreak was confined to the one herd. That herd was depopulated in October, 2008 and the state has maintained its class-free status. Over 8,000 cattle tested negative as part of the mandatory surveillance required before regaining the state's status. Brucellosis vaccination is required statewide with more stringent testing requirements required within the designated surveillance area (DSA). There is official identification of female cattle over 12 months of age statewide, and serologic testing required within 30 days prior to change of ownership, movement from the DSA, interstate movement, or exit from feeder channels. If a cow is tested during the lower-risk period (July 1 – November 1), it can be moved within 60 days. Cattle tested during the higher-risk period (November 2 – June 30) can be moved within 30 days.

Brucellosis Management

The Interagency Brucellosis Management Plan (IBMP) was established in 2000 to manage the risk of *B. abortus* transmission from bison to cattle by implementing hazing, test-and-slaughter, hunting, and other actions near the boundary of Yellowstone National Park (Plumb and Aune, 2002; Donch et al., 2005). To date, these actions have successfully prevented the transmission of *B. abortus* from bison to cattle (Clarke et al., 2005), and an assessment suggested that the risk of future *B. abortus* transmission is minimal under current management conditions (Kilpatrick et al., 2009). Since 2000, about 3200 bison have been removed from the Yellowstone herd with over 1000 animals, or 20% of the total population, culled during the winter of 2005-2006. These actions have been controversial with animal advocacy groups.

The IMBP was not intended to incorporate potential *B. abortus* transmission between elk and bison, and the resultant risks of transmission between elk and cattle. All recent detections of brucellosis in northern GYA cattle have been qualitatively attributed to elk that may or may not have seasonally occupied YNP (Galey et al., 2005). Due to the intense focus on bison *B. abortus* management during the past decade, elk have received minimal brucellosis management attention until recently and often move freely across the ecosystem and come into close contact with cattle premises.

Due to increased *B. abortus* prevalence in Wyoming elk, more elk *B. abortus* mitigation strategies have been evaluated. A five-year pilot test-and-slaughter program in the Pinedale area by Laura Linn-Meadows lowered *B. abortus* seroprevalence but at a cost of \$7000 for each elk removed (USAHA, 2009). The study showed that only 50% of all test-eligible elk were able to be captured. Of all animals sent to slaughter, half were culture-positive.

Since the early period of the BEP, vaccination has been considered a control method for *B. abortus* transmission. The immune-response necessary for conferring protection to the bacteria is a cell-mediated response, especially from CD8 cytotoxic T cells, which is necessary for attacking intracellular bacteria (Schurig et al., 2002). With *B. abortus*, the lipopolysaccharide (LPS) is the major inducer of antibody responses and involves an incompletely understood intracellular signal involving tumor necrosis factor α , perforin, and gamma-interferon (Zhan et al., 1993; Murphy et al., 2001).

There are two USDA-licensed vaccines that offer a measure of protection against *B. abortus*. The strain 19 vaccine was developed from an isolate of *B. abortus* taken from the milk of a Jersey cow in 1923. The isolate was kept at room temperature for over a year and discovered to have lost some of its virulence (Buck, 1930). Unfortunately, the bacterium kept its O side chain of the LPS, which causes animals vaccinated with strain 19 (S19) to test positive on standard *B. abortus* serologic tests. There are also side effects to S19 vaccination. About 1-2.5% of pregnant cattle vaccinated with S19 abort their calves (Manthei, 1952; Mingle, 1961; Beckett and McDiarmid, 1985). There have also been reports and experimental evidence of a more rare association between lameness in cattle and vaccination with strain 19 (Bracewell and Corbel, 1980; Wyn-Jones et al., 1980; Nicoletti et al., 1986; Rogerson and Morgan, 1986; Corbel et al., 1989; Johnson et al., 1994). The arthropathy is caused by *Brucella* antigen-containing immune complexes which locate in the affected joints. Despite the risks of vaccination, elk are vaccinated with S19 on the Wyoming feedgrounds, except for one site for comparison.

Because the serologic cross-reactions of S19 make it ineffective for test-and-cull methods of *B. abortus* control, other candidate vaccines were explored. A live rifampin-resistant “rough”, or devoid of the LPS O-chain, attenuated strain of *B. abortus* labeled “51” by internal laboratory

nomenclature was developed by Schurig and colleagues (1991) and was later trademarked by Virginia Tech Intellectual Properties in 1992. Rough *Brucella* 51 (RB51) has proven to be less abortigenic in cattle than S19 while showing similar efficacy. Because RB51 lacks the O-chain on its LPS it does not cross-react on *B. abortus* serologic tests. The vaccine was licensed for use in cattle by USDA-Animal and Plant Health Inspection Service (APHIS) in 1996. However, it has not shown efficacy in elk, which is why S19 is the only vaccine used in this species (Kreeger et al., 2002).

The efficacy of RB51 in bison remains in dispute (Davis and Elzer, 2002). Olsen et al. (2003) reported that RB51 was efficacious as a calfhood vaccine, whereas data reported by Elzer and Davis (2002) were contradictory. However, there does appear to be consensus that RB51 vaccine is safe as a calfhood vaccine for bison. Also, information about effects on individual bison from vaccination during pregnancy is limited, and there are concerns about abortigenic responses in bison. No abortions occurred when pregnant female bison were vaccinated during their first or second trimesters of gestation (2-5 months after conception; Elzer et al. 1998, Davis and Elzer 2002, Olsen and Holland 2003). However, 2 of 8 pregnant females that were vaccinated during the second half of gestation (i.e., 4.5-6.5 months after conception) aborted their fetuses (Palmer et al. 1996). Also, 50% of seronegative, female bison vaccinated in late pregnancy seroconverted and, while no abortions occurred, both RB51 and field strain *B. abortus* were shed at parturition (Roffe and Olsen 2002).

Age-specific seroprevalence proportions in Yellowstone bison indicate that approximately 50% of bison are exposed prior to reproductive maturity (Treanor et al. 2007). Thus, early exposure to the vaccine may allow immature bison to develop resistance to infection, which could be maintained by booster vaccinations to reduce the occurrence of *B. abortus*-

induced abortions. Seeking to increase tolerance for bison outside YNP and reduce risk of cattle *B. abortus* exposure, the National Park Service has been exploring the option for the remote delivery of the RB51 brucellosis vaccine to various segments of the YNP bison herd (USDI-NPS, 2010). Vaccination of all female bison within YNP is expected to significantly reduce the population seroprevalence of brucellosis (Treanor et al., 2010).

The objective of the vaccination program is to reduce the risk of *B. abortus* transmission to livestock outside YNP by decreasing brucellosis infection in the Yellowstone bison herd. An individual-based model (IBM) was constructed to capture the variability between individuals and estimate responses to both the disease and vaccination for the overall bison population (Treanor et al., 2010). The IBM tracked information on each female bison born into the population. The model used a yearly time step to simulate population level processes and daily time steps to simulate exposure routes during the transmission period. The yearly time step components involved mating, natural mortality, exposure to *B. abortus* via elk, and effects of NPS management operations (testing and subsequent removal of seropositive bison at park boundaries). The daily time step detailed the processes (*B. abortus*-induced abortions and infectious live births) leading to shedding and transmission of *B. abortus* among Yellowstone bison. Demographic, life history, and management-related information (age, sex, disease status, reproductive status, vaccination status, and management removal) were recorded for each female bison modeled.

The estimated brucellosis seroprevalence has fluctuated between 40-60% in YNP bison during the past 20 years (Cheville et al., 1998). This range of infection prevalence was simulated by the model prior to the analysis of each vaccination alternative. The following three vaccination alternatives were simulated: 1) vaccinating female calves and yearlings captured

during boundary management operations, 2) combining remote vaccination using biobullet delivery (Olsen et al., 2006) with boundary vaccination of female calves and yearlings, and 3) vaccinating all female bison during boundary operations as well as by remote delivery. Under each alternative, bison captured at the boundary were tested and test-positive animals removed. The effects of vaccination are likely to play out over a 10-30 year time horizon, during which other ecological factors such as variations in snow pack and predation risk may obscure the effects of vaccination. For example, Cross et al. (2007) suggested winter severity could affect the duration of aggregation by ungulates. If this aggregation coincided with the peak *B. abortus* transmission period, these factors could play an important role in the maintenance of the disease.

Modeling and Risk Assessment

The Yellowstone bison population has been extensively modeled (Peterson et al., 1991a, b; Dobson and Meagher, 1996; Gross et al., 2002; Treanor et al., 2010). However, none of these models attempt to estimate the contact rates required to maintain *B. abortus* at documented prevalence levels. Early modeling studies of *B. abortus* transmission focused exclusively on bison and used mathematical models to show how individual populations interact with a parasitic bacterial pathogen (Peterson et al., 1991a, b; Dobson and Meagher, 1996). Dobson and Meagher's deterministic state-transition model showed that the proportion of hosts infected with *B. abortus* increases as a function of herd density and that 200 infected individuals were necessary to establish *B. abortus* in the Yellowstone bison herd. Once established the authors speculated that high levels of culling would be required to eradicate the disease.

None of the previous modeling efforts attempted to quantify the *B. abortus* transmission pathways within and between bison and elk in the GYA and the risk to cattle from both wildlife

hosts. After decades of high level management and scrutiny, Kilpatrick et al. (2009) provided the first quantitative assessment of the risk of *B. abortus* transmission from Yellowstone bison to cattle grazing in the northern portion of the GYA. Their estimates of bacterial transmission risk were heterogeneous across the spatial landscape and varied with bison population numbers and winter severity. However, Kilpatrick et al. (2009) did not include elk in their analyses or examine explicit spatial information on range overlap between wildlife and cattle.

LITERATURE CITED

2002. Agricultural Bioterrorism Protection Act of 2002; Listing of Biological Agents and Toxins and Requirements and Procedures for Notification of Possession. Department of Agriculture, Animal and Plant Health Inspection Service. RIN 0579-AB47.
- ARENAS, G. N., A. S. STASKEVICH, A. ABALLAY, AND L. S. MAYORGA. 2000. Intracellular trafficking of *Brucella abortus* in J774 macrophages. *Infection and Immunity* 68: 4255-4263.
- AUNE, K., K. ALT, AND T. LEMKE. 2002. Managing wildlife habitat to control brucellosis in the Montana portion of the greater Yellowstone area. . *In* Brucellosis in elk and bison in the greater Yellowstone area. T. J. Kreeger, (ed.). Wyoming Game and Fish Department, Cheyenne, Wyoming, USA. pp.
- AUNE, K., J. C. RHYAN, B. CORSO, AND T. ROFFE. 2007. Environmental persistence of *Brucella* organisms in natural environments of the greater Yellowstone area - A preliminary analysis. Report of the Committee on Brucellosis. Proceedings of the 110th Annual Meeting of the United States Animal Health Association, United States Animal Health Association, Richmond, Virginia.
- BANG, B. 1897. The etiology of epizootic abortion. *Journal of Comparative Pathology and Therapeutics* 10: 125-149.
- BECKETT, F. W., AND S. C. MCDIARMID. 1985. The effect of reduced-dose *Brucella abortus* strain 19 vaccination in accredited dairy herds. *British Veterinary Journal* 141: 507.

- BELLOU, L., M. JANOVSKY, AND E. M. VILEI, ET AL. 2004. Molecular epidemiology of *Mycoplasma conjunctivae* in caprinae: Transmission across species in natural outbreaks. *Applied and Environmental Microbiology* 69: 1913-1919.
- BERCOVICH, Z. 1998. Maintenance of *Brucella abortus*-free herds: A review with emphasis on the epidemiology and the problems in diagnosing brucellosis in areas of low prevalence. *Veterinary Quarterly* 20: 81-88.
- BERGER, J., AND S. L. CAIN. 1999. Reproductive synchrony in brucellosis-exposed bison in the southern greater Yellowstone ecosystem and in noninfected populations. *Conservation Biology* 13: 357-366.
- BITTNER, A. 2004. An overview and the economic impacts associated with mandatory brucellosis testing in Wyoming cattle. Wyoming Department of Administration and Information, Economic Analysis Division.
- BÖHM, M., M. R. HUTCHINGS, AND P. C. L. WHITE. 2009. Contact networks in a wildlife-livestock host community: Identifying high-risk individuals in the transmission of bovine TB among badgers and cattle. *PLoS ONE* 4: e5016.
- BRACEWELL, C. D., AND M. J. CORBEL. 1980. An association between arthritis and persistent serological reactions to *Brucella abortus* in cattle from apparently brucellosis-free herds. *Veterinary Record* 106.
- BUCK, J. M. 1930. Studies of vaccination during calfhood to prevent bovine infectious abortion. *Journal of Agricultural Research* 41: 667.
- CATLIN, J. E., AND E. J. SHEEHAN. 1986. Transmission of bovine brucellosis from dam to offspring. *Journal of the American Veterinary Medical Association* 188: 867-869.

- CENTERS FOR DISEASE CONTROL AND PREVENTION. 2007. Disease listing, Brucellosis, General Information.
http://www.cdc.gov/ncidod/dbmd/diseaseinfo/brucellosis_g.htm#ivet.
- CHEVILLE, N., D. R. MCCULLOUGH, AND L. R. PAULSON. 1998. Brucellosis in the greater Yellowstone area. National Research Council, Washington D.C.
- CHIVIAN, E. 2001. Environment and health: 7. Species loss and ecosystem disruption-the implications for human health. *Journal of the Canadian Medical Association* 164: 66-69.
- CHOMEL, B. B., E. E. DEBESS, D. M. MANGIAMELE, K. F. REILLY, T. B. FARVER, R. K. SUN, AND L. R. BARRETT. 1994. Changing trends in the epidemiology of human brucellosis in California from 1973 to 1992: a shift toward foodborne transmission. *Journal of Infectious Diseases* 170: 1216-1223.
- CLARKE, R., C. JOURDONNAIS, J. MUNDINGER, L. STOEFFLER, AND R. WALLEN. 2005. A Status Review of Adaptive Management Elements 2000 to 2005. Interagency Bison Management Plan.
- CLIFFORD, D. L., B. A. SCHUMAKER, T. R. STEPHENSON, V. C. BLEICH, M. L. CAHN, B. J. GONZALES, W. M. BOYCE, AND J. A. K. MAZET. 2009. Assessing disease risk at the wildlife-livestock interface: A study of Sierra Nevada bighorn sheep. *Biological Conservation* 142: 2559-2568.
- COE, P. K., B. K. JOHNSON, K. M. STEWART, AND J. G. KIE. 2005. Spatial and temporal interactions of elk, mule deer and cattle. *In* The Starkey Project: a synthesis of long-term studies of elk and mule deer. Reprinted from the 2004 Transactions of the North American Wildlife and Natural Resources Conference. M. J. Wisdom, (ed.). Alliance Communications Group, Lawrence, Kansas, USA. pp. 150-158.

- COOK, W. E., E. S. WILLIAMS, AND S. A. DUBAY. 2004. Disappearance of bovine fetuses in northwestern Wyoming. *Wildlife Society Bulletin* 32: 254-259.
- CORBEL, M. J., F. A. STUART, AND R. A. BREWER, ET AL. 1989. Arthropathy associated with *Brucella abortus* strain 19 vaccination in cattle. I. Examination of field cases. *British Veterinary Journal* 145: 337-346.
- CREECH, G. T. 1930. *Brucella abortus* infection in a male bison. *North American Veterinarian* 11: 35-36.
- CROSS, P. C., W. H. EDWARDS, B. M. SCURLOCK, E. J. MAICHAK, AND J. D. ROGERSON. 2007. Effects of management and climate on elk brucellosis in the Greater Yellowstone Ecosystem. *Ecological Applications* 17: 957-964.
- CROSS, P. C., T. O. LEMKE, P. J. WHITE, AND D. B. TYERS. 2009. Northern Yellowstone cooperative wildlife working group 2008 annual report (October 1, 2007-September 30, 2008). U.S. Geological Survey, Northern Rocky Mountain Science Center, Bozeman, Montana.
- CUNNINGHAM, A. A. 2005. A walk on the wild side - emerging wildlife diseases: They increasingly threaten human and animal health. *British Medical Journal* 331: 1214-1215.
- DASZAK, P., A. A. CUNNINGHAM, AND A. D. HYATT. 2000. Emerging infectious diseases of wildlife -- Threats to biodiversity and human health. *Science* 287: 443-449.
- DASZAK, P., A. A. CUNNINGHAM, AND A. D. HYATT. 2001. Anthropogenic environmental change and the emergence of infectious diseases in wildlife. *Acta Tropica*: 103-116.

- DAVIS, D., J. W. TEMPLETON, T. A. FICHT, J. D. WILLIAMS, J. D. KOPEC, AND L. G. ADAMS. 1990. *Brucella abortus* in captive bison. I. Serology, bacteriology, pathogenesis, and transmission to cattle. *Journal of Wildlife Diseases* 26: 360-371.
- DAVIS, D. S., AND P. H. ELZER. 2002. *Brucella* vaccines in wildlife. *Veterinary Microbiology* 90: 533-544.
- DAVIS, D. S., J. W. TEMPLETON, T. A. FICHT, J. D. HUBER, R. D. ANGUS, AND L. G. ADAMS. 1991. *Brucella abortus* in bison. II. Evaluation of strain 19 vaccination of pregnant cows *Journal of Wildlife Diseases* 27: 258-264.
- DELAHAY, R. J., G. C. SMITH, AND M. R. HUTCHINGS. 2009. The Science of Wildlife Disease Management. *In* Management of Disease in Wild Mammals. R. J. Delahay, G. C. Smith, AND M. R. Hutchings, (eds.). Spring, Tokyo, Japan. pp. 1-8.
- DICKSON, T. 2005. Taking aim at deprecation; Montana's new liberalized elk and deer season is designed to help landowners and hunters. *Montana Outdoors*. Montana Fish, Wildlife, and Parks.
- DOBSON, A. 2004. Population Dynamics of Pathogens with Multiple Host Species. *The American Naturalist* 164: S64-S78.
- DOBSON, A., AND M. MEAGHER. 1996. The Population Dynamics of Brucellosis in the Yellowstone National Park. *Ecology* 77: 1026-1036.
- DONCH, D. A., AND A. A. GERTONSON. 2008. Status report -- Fiscal year 2008; Cooperative State-Federal Brucellosis Eradication Program. USDA-APHIS Veterinary Services.
- DONCH, D. A., A. A. GERTONSON, J. C. RHYAN, AND M. J. GILSDORF. 2005. Status Report - Fiscal Year 2005; Cooperative State-Federal Brucellosis Eradication Program. USDA-APHIS Veterinary Services.

- EBERHARDT, L. L., P. J. WHITE, R. A. GARROTT, AND D. B. HOUSTON. 2007. A seventy-year history of trends in Yellowstone's northern elk herd. *Journal of Wildlife Management* 71: 594-602.
- EDMONDS, M. D., F. M. WARD, T. M. O'HARA, AND P. H. ELZER. 1999. Use of western immunoblot analysis for testing moose serum for *Brucella suis* biovar 4 specific antibodies. *Journal of Wildlife Disease* 35: 591-595.
- ETTER, R. P., AND M. L. DREW. 2006. Brucellosis in Elk of Eastern Idaho. *Journal of Wildlife Diseases* 42: 271-278.
- FERRARI, M. J., AND R. A. GARROT. 2002. Bison and Elk: Brucellosis Seroprevalence on a Shared Winter Range. *The Journal of Wildlife Management* 66: 1236-1254.
- FLAGG, D. E. 1983. A case history of a brucellosis outbreak in a brucellosis free state which originated in bison. *In Proceedings: Proceedings of the United States Animal Health Association*. pp. 171-172.
- FRENCHICK, P. J., R. J. MARKHAM, AND A. H. COCHRANE. 1985. Inhibition of phagosome-lysosome fusion in macrophages by soluble extracts of virulent *Brucella abortus*. *American Journal of Veterinary Research* 46: 332-335.
- FULLER, J. A., R. A. GARROTT, P. J. WHITE, K. E. AUNE, T. J. ROFFE, AND J. C. RHYAN. 2007. Reproduction and survival of Yellowstone bison. *Journal of Wildlife Management* 71: 2365-2372.
- GALEY, F., J. BOUSMAN, T. CLEVELAND, J. ETCHPARE, R. HENDRY, AND J. HINES, ET AL. 2005. Wyoming Brucellosis Coordination Team report and recommendations. Report presented to Governor Dave Freudenthal. Cheyenne, Wyoming, USA.

- GALL, D., AND K. NIELSEN. 2004. Serological diagnosis of bovine brucellosis: A review of test performance and cost comparison. *Revue Scientifique et Technique Office International des Epizooties* 23: 989-1002.
- GALL, D., K. NIELSEN, L. FORDES, D. DAVIS, P. H. ELZER, S. C. OLSEN, S. BALSEVICIUS, L. KELLY, P. SMITH, S. TAN, AND D. JOLY. 2000. Validation of the fluorescence polarization assay and comparison to other serological assays for the detection of serum antibodies to *Brucella abortus* in bison. *Journal of Wildlife Diseases* 36: 469-476.
- GODFROID, J. 2002. Brucellosis in wildlife. *Revue Scientifique et Technique Office International des Epizooties* 21: 277-286.
- GOVERNMENT ACCOUNTABILITY OFFICE. 2009. Veterinarian Workforce: Actions are Needed to Ensure Sufficient Capacity for Protecting Public and Animal Health. GAO-09-178, Government Accountability Office, Washington, D.C.
- GROSS, J. E., B. C. LUBOW, AND M. W. MILLER. 2002. Modeling the epidemiology of brucellosis in the Greater Yellowstone Area. *In* Brucellosis in elk and bison in the Greater Yellowstone Area. T. J. Kreeger, (ed.). Wyoming Game and Fish Dept., Cheyenne, WY. pp. 24-37.
- HALBERT, N. D., T. J. WARD, R. D. SCHNABEL, J. F. TAYLOR, AND J. N. DERR. 2005. Conservation genomics: Disequilibrium mapping of domestic cattle chromosomal segments in North American bison populations. *Molecular Ecology* 14: 2343-2362.
- HAMLIN, K. L., AND J. A. CUNNINGHAM. 2008. Montana elk movements, distribution, and numbers relative to brucellosis transmission risk. *Montana Fish, Wildlife, and Parks*.

- JOHNSON, B., D. A. MOSIER, R. J. MORTON, AND A. W. CONFER. 1994. Experimental *Brucella abortus* strain 19 arthritis in young cattle. *Journal of Veterinary Diagnostic Investigation* 6: 56-61.
- JOHNSON, D. E. 1951. Biology of the elk calf, *Cervus canadensis nelsoni*. *Journal of Wildlife Management* 15: 396-410.
- JOLY, D. O., AND F. MESSIER. 2005. The effect of bovine tuberculosis and brucellosis on reproduction and survival of wood bison in Wood Buffalo National Park. *Journal of Animal Ecology* 74: 543-551.
- JONES, J. D., J. J. TREANOR, AND R. L. WALLEN. 2009. Parturition in Yellowstone bison. YCR-2009-01, National Park Service, Yellowstone National Park, Wyoming.
- KILPATRICK, A. M., C. M. GILLIN, AND P. DASZAK. 2009. Wildlife-livestock conflict: the risk of pathogen transmission from bison to cattle outside Yellowstone National Park. *Journal of Applied Ecology* 46: 476-485.
- KITTELBERGER, R., F. HILBANK, M. F. HANSEN, M. PENROSE, G. W. DE LISLE, J. J. LETESSON, B. GARIN-BASTUJI, J. SEARSON, C. A. FOSSATI, A. CLOECKAERT, AND G. G. SCHURIG. 1995. Serological crossreactivity between *Brucella abortus* and *Yersinia enterocolitica* O:9 I immunoblot analysis of the antibody response to *Brucella* protein antigens in bovine brucellosis. *Veterinary Microbiology* 47: 257-270.
- KNOX, W. D. 1947. Control of brucellosis in dairy herds. *In Proceedings: Proceedings of the Fifty-fifth Annual Meeting of the United States Livestock Sanitary Association*. Waverly Press. pp. 144-149.
- KOZUKEEV, T. B., S. AJEILAT, E. MAES, AND M. FAVOROV. 2006. Risk factors for brucellosis -- Leylek and Kadamjay districts, Batken Oblast, Kyrgyzstan, January-

- November, 2003. Morbidity and Mortality Weekly Report, Centers for Disease Control and Prevention, Atlanta, Georgia.
- KREEGER, T. J., W. E. COOK, W. H. EDWARDS, P. H. ELZER, AND S. C. OLSEN. 2002. *Brucella abortus* strain RB51 vaccination in elk II. Failure of high dosage to prevent abortion. *Journal of Wildlife Diseases* 38: 27-31.
- LAPRAIK, R. D., D. D. BROWN, H. MANN, AND T. BRAND. 1975. Brucellosis: a study of five calves from reactor dams. *Veterinary Record* 97: 52-54.
- MAICHAK, E. J., B. M. SCURLOCK, J. D. ROGERSON, L. L. MEADOWS, A. E. BARBKNECHT, W. H. EDWARDS, AND P. C. CROSS. 2009. Effects of management, behavior, and scavenging on risk of brucellosis transmission in elk of western Wyoming. *Journal of Wildlife Diseases* 45: 398-410.
- MANTHEI, C. A. 1952. Evaluation of vaccinal methods and doses of *Brucella abortus* strain 19. *In Proceedings: Proceedings of the 56th Annual Meeting of the Livestock Sanitation Association.* pp. 115.
- MANTHEI, C. A., AND R. W. CARTER. 1950. Persistence of *Brucella abortus* infection in cattle. *American Journal of Veterinary Research* 11: 173-180.
- MEAGHER, M. 1973. The bison of Yellowstone National Park. National Park Service, Washington, D.C., USA.
- MEAGHER, M., AND M. E. MEYER. 1994. On the origin of brucellosis in bison of Yellowstone National Park: A review. *Conservation Biology* 8: 645-653.
- MINGLE, C. K. 1961. Cooperative state-federal brucellosis eradication. *In Proceedings: Proceedings of the Sixty-fifth Annual Meeting of the United States Livestock Sanitary Association.* MacCrellish and Quigley. pp. 108-119.

- MORTON, J. K., E. T. THORNE, AND G. M. THOMAS. 1981. Brucellosis in elk III. Serologic evaluation. *Journal of Wildlife Diseases* 17: 23-31.
- MURPHY, E. A., J. SATHIYASEELAN, M. A. PARENT, B. ZOU, AND C. L. BALDWIN. 2001. Interferon-gamma is crucial for surviving a *Brucella abortus* infection in both resistant C57BL/6 and susceptible BABB/c mice. *Immunology* 103: 511-518.
- NICOLETTI, P., A. M. CROWLEY, J. A. RICHARDSON, AND J. A. FARRAR. 1986. Suspected *Brucella abortus* strain 19-induced arthritis in a dairy cow. *Agri-Practice* 7: 5-6.
- OLMSTEAD, A. L., AND P. W. RHODE. 2004. An Impossible Undertaking: The Eradication of Bovine Tuberculosis in the United States. *The Journal of Economic History* 64: 734-772.
- OLSEN, S., A. E. JENSEN, W. C. STOFFREGEN, AND M. V. PALMER. 2003. Efficacy of calfhood vaccination with *Brucella abortus* strain RB51 in protecting bison against brucellosis. *Research in Veterinary Science* 74: 17-22.
- OLSEN, S. C., R. J. CHRISTIE, D. W. GRAINGER, AND W. S. STOFFREGEN. 2006. Immunologic responses of bison to vaccination with *Brucella abortus* strain RB51: comparison of parenteral to ballistic delivery via compressed pellets or photopolymerized hydrogels. *Vaccine* 24: 1346-1353.
- PAPPAS, G., P. PAPADIMITRIOU, N. AKRITIDIS, L. CHRISTOU, AND E. TSIANOS. 2006. The new global map of human brucellosis. *Lancet Infectious Diseases* 6: 91-99.
- PETERSON, M. J., W. E. GRANT, AND D. S. DAVIS. 1991a. Bison-Brucellosis Management - Simulation of Alternative Strategies. *Journal of Wildlife Management* 55: 205-213.

- PETERSON, M. J., W. E. GRANT, AND D. S. DAVIS. 1991b. Simulation of Host-Parasite Interactions within a Resource-Management Framework - Impact of Brucellosis on Bison Population-Dynamics. *Ecological Modelling* 54: 299-320.
- PIZARRO-CERDA, J., E. MORENO, V. SANGUEDOLCE, J. L. MEGE, AND J. P. GORVEL. 1998. Virulent *Brucella abortus* prevents lysosome fusion and is distributed within autophagosome-like compartments. *Infectious Immunology* 66: 2387-2392.
- PLOMMET, M., R. FENSTERBANK, G. RENOUX, J. GESTIN, AND A. PHILIPPON. 1973. Brucellose bovine experimentale XII - Persistence a l'age adulte de l'infection congenitale de la genisse. *Annals of Veterinary Research* 4: 419-435.
- PLUMB, G., AND K. AUNE. 2002. The long term Interagency Bison Management Plan for Yellowstone National Park and the State of Montana. *In* Brucellosis in elk and bison in the Greater Yellowstone Area. T. J. Kreeger, (ed.). Wyoming Department of Game and Fish, Cheyenne, WY. pp. 136-145.
- PLUMB, G. E., P. J. WHITE, M. B. COUGHENOUR, AND R. L. WALLEN. 2009. Carrying capacity, migration, and dispersal in Yellowstone bison. *Biological Conservation* 142: 2377-2387.
- POWER, A. G., AND C. E. MITCHELL. 2004. Pathogen Spillover in Disease Epidemics. *The American Naturalist* 164: S79-S89.
- PROFFITT, K. M., P. J. WHITE, AND R. A. GARROTT. 2010. Spatio-temporal overlap between Yellowstone bison and elk -- implications of wolf restoration and other factors for brucellosis transmission risk. *Journal of Applied Ecology* 47: 281-289.
- RAGAN, V. E. 2002. The Animal and Plant Health Inspection Service (APHIS) brucellosis eradication program in the United States. *Veterinary Microbiology* 90: 11-18.

- RHYAN, J. C., T. GIDLEWSKI, T. J. ROFFE, K. AUNE, L. M. PHILO, AND D. R. EWALT. 2001. Pathology of brucellosis in bison from Yellowstone National Park. *Journal of Wildlife Diseases* 37: 101-109.
- RHYAN, J. C., S. D. HOLLAND, T. GIDLEWSKI, D. A. SAARI, A. E. JENSEN, D. R. EWALT, S. G. HENNAGER, S. C. OLSEN, AND N. F. CHEVILLE. 1997. Seminal vesiculitis and orchitis caused by *Brucella abortus* biovar 1 in young bison bulls from South Dakota. *Journal of Veterinary Diagnostic Investigation* 9: 368-374.
- ROFFE, T. J., J. C. RHYAN, K. AUNE, L. M. PHILO, D. R. EWALT, T. GIDLEWSKI, AND S. G. HENNAGER. 1999. Brucellosis in Yellowstone National Park bison: Quantitative serology and infection. *Journal of Wildlife Management* 63: 1132-1137.
- ROGERSON, B. A., AND I. R. MORGAN. 1986. Investigation of aberrant positive reactions to serological tests for bovine brucellosis. *Australian Veterinary Journal* 63: 227-229.
- SCHUMAKER, B. A., J. A. K. MAZET, B. J. GONZALES, P. H. ELZER, S. K. HIETALA, AND M. H. ZICCARDI. 2010. Evaluation of the western immunoblot as a detection method for *Brucella abortus* exposure in elk. *Journal of Wildlife Diseases* 46: 87-94.
- SCHURIG, G. G., R. M. ROOP II, T. BAGCHI, S. BOYLE, D. BUHRMAN, AND N. SRIRANGANATHAN. 1991. Biological properties of RB51; a stable rough strain of *Brucella abortus*. *Veterinary Microbiology* 28: 171-188.
- SCHURIG, G. G., N. SRIRANGANATHAN, AND M. J. CORBEL. 2002. Brucellosis vaccines: past, present and future. *Veterinary Microbiology* 90: 479-496.
- SETON, E. T. 1927. *Lives of game animals*. Doubleday, Page and Company, Garden City, New York, USA.

- SMITH, B. L. 2001. Winter feeding of elk in western North America. *Journal of Wildlife Management* 65: 173-190.
- STATE OF MICHIGAN. 2008. Emerging disease issues: Management of bovine tuberculosis in Michigan deer. http://www.michigan.gov/emergingdiseases/0,1607,7-186-25804_25811-75930--,00.html.
- SUTHERLAND, S. S. 1980. Immunology of bovine brucellosis. *Veterinary Bulletin* 50: 359-368.
- TAYLOR, D. T., E. B. BRADLEY, AND M. M. MARTIN. 1981. The outfitting industry in Teton County: its clientele and economic importance. Agricultural Extension Service Publication B-793, University of Wyoming, Laramie, WY, USA.
- TAYLOR, L. H., S. M. LATHAM, AND M. E. J. WOOLHOUSE. 2001. Risk factors for human disease emergence. *Philosophical Transactions of the Royal Society of London* 356: 983-989.
- THORNE, E. T. 2001. Brucellosis. *In Infectious Diseases of Wild Mammals*. E. S. Williams, AND I. K. Barker, (eds.). Blackwell Publishing, Ames, Iowa. pp. 372-395.
- THORNE, E. T., J. K. MORTON, F. M. BLUNT, AND H. A. DAWSON. 1978a. Brucellosis in elk. II. Clinical effects and means of transmission as determined through artificial infections. *Journal of Wildlife Diseases* 14: 280-291.
- THORNE, E. T., J. K. MORTON, AND W. C. RAY. 1979. Brucellosis, its effect and impact on elk in western Wyoming. *In North American elk: ecology, behavior and management*. M. S. Boyce, AND L. D. Hayden-Wing, (eds.). The University of Wyoming, Laramie, WY. pp. 212-220.

- THORNE, E. T., J. K. MORTON, AND G. M. THOMAS. 1978b. Brucellosis in elk I. Serologic and bacteriologic survey in Wyoming. *Journal of Wildlife Diseases* 14: 74-81.
- TOMLEY, F. M., AND M. W. SHIRLEY. 2009. Livestock infectious diseases and zoonoses. *Philosophical Transactions of the Royal Society Series B* 364: 2637-2642.
- TREANOR, J., J. JOHNSON, R. WALLEN, S. CILLES, P. CROWLEY, AND D. MAEHR. 2008. Vaccination strategies for managing brucellosis in Yellowstone bison. YCR-2008-03, National Park Service, Yellowstone National Park, Wyoming.
- TREANOR, J. J., J. S. JOHNSON, R. L. WALLEN, S. CILLES, P. H. CROWLEY, J. J. COX, D. S. MAEHR, P. J. WHITE, AND G. E. PLUMB. 2010. Vaccination strategies for managing brucellosis in Yellowstone bison. *Vaccine* 28S: F64-F72.
- UN OFFICE FOR THE COORDINATION OF HUMANITARIAN AFFAIRS. 2003. Kyrgyzstan: Focus on brucellosis in south. IRIN humanitarian news and analysis, Osh.
- USAHA. 2009. Report of the Committee on Brucellosis. United States Animal Health Association, San Diego, CA.
- USDA FOREST SERVICE, I. R. 2010. Southwest Idaho Ecogroup Land and Resource Management Plans, Final Supplemental Environmental Impact Statement.
- USDI-NPS. 2010. Draft Environmental Impact Statement for Brucellosis Remote Vaccination Program for Bison in Yellowstone National Park. US Department of Interior.
- VAN DEN BORN, J. M., AND D. J. VERVOORN. 1965. Control of bovine tuberculosis and brucellosis in the Netherlands. *Bulletin-Office International des Epizooties* 63: 1531-1554.
- WESTERN, D. 2001. Human-modified ecosystems and future evolution. *Proceedings of the National Academy of Science* 98: 5458-5465.

- WILESMITH, J. W. 1978. The persistence of *Brucella abortus* infection in calves: a retrospective study of heavily infected herds. *Veterinary Record* 103: 149-153.
- WILSON, D. W., AND P. T. BEERS. 2001. Global trade requirements and compliance with World Trade Organization agreements: the role of tracing animals and animal products. *Revue Scientifique et Technique Office International des Epizooties* 20: 379-384.
- WISE, R. I. 1980. Brucellosis in the United States: Past, present, and future. *Journal of the American Medical Association* 244: 2318-2322.
- WOOLHOUSE, M. E. J., L. H. TAYLOER, AND D. T. HAYDON. 2001. Population Biology of Multihost Pathogens. *Science* 292: 1109-1112.
- WYN-JONES, G., J. R. BAKER, AND P. M. JOHNSON. 1980. A clinical and immunopathological study of *Brucella abortus* strain 19-induced arthritis in cattle. *Veterinary Record* 107: 5-9.
- YOUNG, E. J., AND U. SUVANNOPARRAT. 1975. Brucellosis outbreak attributed to ingestion of unpasteurized goat cheese. *Archives of Internal Medicine* 135: 240-243.
- ZHAN, Y., J. CHANG, AND C. CHEERS. 1993. Cytokine response of T-cell subsets from *Brucella abortus* infected mice to soluble *Brucella* proteins. *Infectious Immunology* 61: 2841-2847.

Objective 1

BISON OR ELK: WHO SHOULD BE THE TARGET OF BRUCELLOSIS CONTROL IN THE NORTHERN GREATER YELLOWSTONE AREA?

Brant A. Schumaker,^{1,2*} Jonna A.K. Mazet,² John Treanor,³ Rick Wallen,³ Anthony Tam,¹ Ian A. Gardner,^{2,4} Martin Zaluski,⁵ and Tim Carpenter^{1,2}

¹Center for Animal Disease Modeling and Surveillance (CADMS), Department of Medicine and Epidemiology, School of Veterinary Medicine, University of California, Davis, CA 95616, USA;

²Wildlife Health Center, One Health Institute, School of Veterinary Medicine, University of California, Davis, CA 95616, USA; ³National Park Service, Yellowstone National Park, P.O. Box 168, Wyoming 82190, USA; ⁴Department of Medicine and Epidemiology, School of Veterinary Medicine, University of California, Davis, CA 95616, USA; ⁵Montana Department of Livestock, P.O. Box 202001, Helena, MT 59620-2001, USA;

*Corresponding author (email: theschu@ucdavis.edu, phone: (530) 752-3566, fax: (530) 752-1618)

Abstract

Wild, free-ranging, bison and elk in the greater Yellowstone area (GYA) are the last reported alternate hosts of *Brucella abortus*-caused brucellosis in the United States. The ability of bison and elk to concomitantly serve as reservoirs of *B. abortus* increases the complexity of risk of transmission to cattle, and presents serious challenges for comprehensive disease management. We present the first spatially-explicit risk assessment of brucellosis transmission among elk, bison, and cattle in the northern portion of the GYA. We used a modeling approach based on spatio-temporal probabilities of bacterial shedding by bison and elk on the northern GYA landscape. Interactive effects between population size and winter severity were major determinants influencing bison movements to lower elevation winter grazing areas, overlapping with federally-regulated domestic cattle grazing allotments. Increasing population size resulted in higher herd densities and increased bacterial shedding. Median total risk to cattle from elk and bison was 3.6 cattle-exposure event-days (95% P.I. 0.1-36.6). The estimated percentage of cattle exposure risk from the Yellowstone bison herd was small (0.0-0.3% of total risk) compared with elk which contributed 99.7-100% of the total risk. Natural herd migration and boundary management operations were important in minimizing the contribution of bison to cattle exposure risk, which supports continued boundary management operations for spatio-temporal separation between bison and cattle. Under current management practices, bison risk to cattle grazing in the northern portion of the GYA is expected to be minimal. The comingling of cattle and elk, especially during the late gestation period for elk, should be reduced, as spontaneous elk abortions pose a risk for interspecies disease transmission.

Keywords: bison, brucellosis, disease management, disease modeling, elk, GIS, population dynamics, risk model, spatially explicit model, wildlife health

INTRODUCTION

Wild, free-ranging, bison and elk in the greater Yellowstone area (GYA) are the last-known reservoirs of bovine brucellosis (*Brucella abortus*) in the United States (Godfroid 2002). Because both bison and elk are competent reservoirs of *B. abortus*, comprehensive disease management is a challenge and the aggregated risks of pathogen transmission to cattle is increasingly complicated (Delahay et al. 2009). In addition to *B. abortus*, many multiple-host pathogens cause the most important livestock diseases listed by the World Organisation for Animal Health, including rinderpest, foot-and-mouth disease, Johne's disease, and Rift Valley fever and comprise 80% of the pathogens of domestic animals (Woolhouse et al. 2001).

Understanding the interspecies transmission dynamics of a multi-host system is crucial for disease management (Dobson 2004, Delahay et al. 2009). Some hosts may be persistent reservoirs of pathogens and others may be recurrently infected through spillover (Power and Mitchell 2004). In addition, multi-host systems present differing and complex surveillance and control challenges. Effective surveillance plans provide early detection of emerging infectious diseases and "spillover" disease events, provided the diagnostic tests used are accurate. However, when using diagnostic tests on species for which the test was not developed, there are many issues with test performance. In addition, there can be problems with cross-reactivity on diagnostic tests by commensal organisms of no disease significance (Kittelberger et al. 1995). Overall, having multiple wildlife hosts and reservoirs greatly complicates disease management (Government Accountability Office 2009).

Brucellosis was first detected in Yellowstone bison in 1917 (Mohler 1917) and was most likely introduced from domestic cattle (Meagher and Meyer 1994). The control and eradication of the disease from the United States has been a priority since 1934, when the federal

government sought to reduce the prevalence of the most significant livestock disease at that time. At different times and under different jurisdictions, brucellosis management strategies have included combinations of capture, test, and slaughter of test-positive animals; vaccination; surveillance; and forced spatial-temporal separation from livestock through hazing or slaughter (Donch et al. 2005). Since the inception of the Interagency Bison Management Plan in 2000, bison have been actively managed to prevent spatio-temporal overlap with cattle (Clarke et al. 2005). This active management of bison has prevented bison-cattle transmission of *Brucella*; however, until recently, elk have received minimal risk management attention. In the last decade, there have been multiple detections of brucellosis in cattle in the GYA states (Idaho, Montana, Wyoming), with elk identified as the source of infection for nine cases since 2002 (Donch and Gertonson 2008). Bison conservation continues to be a priority of the National Park Service; however, for decades, livestock and regulatory personnel have viewed Yellowstone bison as the highest priority wildlife source of transmission of pathogens for livestock in the GYA (Meagher and Meyer 1994).

Kilpatrick et al. (2009) provided the first quantitative assessment of the risk of *B. abortus* transmission from Yellowstone bison to cattle grazing in the northern portion of the GYA. Their estimates of bacterial transmission risk were heterogeneous across the spatial landscape and varied with bison population numbers and winter severity. We extended this work by (1) including elk as a source of *B. abortus* transmission, (2) evaluating explicit spatial information on range overlap between wildlife and cattle, (3) providing risk estimates of *Brucella* transmission from both bison and elk to cattle grazing in southwestern Montana, and (4) evaluating the role of winter severity and population size on the spatial distributions of bison and elk and, hence, the overall potential for brucellosis transmission to cattle.

MATERIALS AND METHODS

Study area and wildlife host populations

The GYA is one of the largest intact temperate zone ecosystems on earth and includes portions of Wyoming, Idaho, and Montana. It is also home to the largest wild and free-ranging elk and bison populations in the United States. Elk and bison populations in the GYA are variably infected with *B. abortus*, the cause of cattle brucellosis. Elk populations in northern GYA have a low seroprevalence (i.e., exposure; <5%) of *B. abortus*, whereas seroprevalence in Yellowstone bison is high (40-70%) (Cheville et al. 1998).

One bison population with between 2000 and 5000 individuals (Meagher 1973, Clarke et al. 2005) and five elk populations (Gallatin-Madison, Gravelly-Snowcrest, Madison-Firehole, northern Yellowstone, and Sand Creek, Idaho) are distributed across 3,000 km² in the northern GYA. Estimates of northern Yellowstone elk were near 25,000 animals in the late 1980s, but decreased by approximately 50-60% by 2006 (Eberhardt et al. 2007). Domestic cattle (266 in the winter and 1363 in the spring in 2006) are grazed on public and private lands adjacent to Yellowstone National Park (YNP) and within habitat occupied by bison and elk during the winter (Kilpatrick et al. 2009). Federal and state management agencies have attempted to decrease the risk of *B. abortus* transmission from bison to cattle using hazing and bison culling to maintain spatio-temporal separation from cattle (U.S. Department of Interior [USDI] and U.S. Department of Agriculture [USDA] 2000).

Brucellosis infection and transmission

For *B. abortus* transmission to occur from wildlife to cattle, the following requirements must be met: (1) the wildlife must be infected; (2) infected wildlife must be on allotments or private land where cattle are grazed outside of the National Park; (3) pregnant wildlife must shed *Brucella* into environment (through abortion, birth fluids, or post-partum via placenta); and (4) *B. abortus* must persist on the landscape long enough for grazing cattle to come into contact with bacteria. The probability of *B. abortus* transmission between elk (or from elk to cattle) is likely low during calving (May through June) because pregnant dams isolate themselves while giving birth and meticulously clean the birth site (Johnson 1951). Thus, birth sites are dispersed, and the likelihood of other elk encountering infected birth tissues and fluids is low. However, transmission risk is likely higher during the brucellosis abortion period from February through April when many elk aggregate in larger groups on lower-elevation winter ranges that sometimes include ranch areas with cattle (Hamlin and Cunningham 2008).

Risk model development

We assessed the risk of bacterial shedding from third-trimester abortions and infectious live parturition events for bison and elk populations in the northern GYA. We created a stochastic *Brucella* shedding risk model, which was parameterized using a combination of published peer-reviewed data, unpublished data, and expert opinion on winter severity, animal locations, serologic test results, and population demography (Table 1, also see Supporting Information). We fit statistical distributions to data using @RISK v5.0 (Palisade Corporation, Ithaca, New York) to address the variability and uncertainty of parameters.

Exposure area and wildlife tolerance

It is unknown how close a susceptible cow would have to be to *B. abortus*-infected tissues before it would be likely to investigate them and become exposed. The Starkey Project at the Pacific Northwest Research Station found that forage competition between elk and cattle likely decreases the chance of comingling on winter range (Coe et al. 2005). However, lack of available forage and other environmental pressures during severe winters in the GYA likely increase comingling which is observed annually by National Park Service staff. Because there was a high degree of uncertainty associated with this comingling parameter, we chose to model it as a discrete distribution with equal probabilities from 50 to 250 meters, by 50-meter increments. We then evaluated the effect of this assumption in a sensitivity analysis.

To account for active management of bison in contrast to elk, we included a wildlife tolerance factor. The tolerance factor was defined as how much access bison are given to cattle grazing allotments as a percentage of the access given to elk. Because there are no specific data on this parameter, we modeled it as a uniform distribution between 0-100% and evaluated it in the sensitivity analysis with 10% increments from 0-100%.

Birth synchrony

In general, data suggest bison and elk in the northern portion of the GYA exhibit a high degree of birth synchrony, with the majority (80%) of bison calving during late-April to late-May and elk calving between mid-May to mid-June (Cheville et al. 1998, Berger and Cain 1999). Feed ground data from the southern portion of the GYA in Wyoming have shown birth dates for elk later in the year, but parturition events are still unlikely after the third week of June due to the normal pattern of sexual segregation (Cross et al. 2009, Maichak et al. 2009). We assumed a 285-day gestation period for bison and a 250-day gestation period for elk, with the

initiation of an abortion window for bison in January and for elk in the second week of February (Fig. 1). The model parameterization is consistent with the timing of culture-positive results from aborted elk fetuses submitted by personnel from the Wyoming Game and Fish Department (Cross et al. 2009, Maichak et al. 2009). We fit statistical distributions to parturition data obtained from published and unpublished sources and used our risk model to estimate the percentage of pregnancies that would fail or result in a live parturition with the potential for bacterial shedding using @RISK v5.0 (Palisade Corporation, Ithaca, New York).

Bacterial versus fetal tissue persistence

The number of days a *Brucella*-contaminated birth site is infective is dependent upon the amount of time that it takes for an infected fetus or tissues to be scavenged or for ultraviolet radiation to degrade the bacteria. Aune et al. (2007); unpublished data) and Cook et al. (2004) found that fetuses would be scavenged prior to ultraviolet degradation of bacteria (mean \pm SD = 18.2 ± 20.1 days). We used a distribution with similar characteristics, BetaGeneral(2, 6.93, 1, 78), for consistency and comparability with other models (Kilpatrick et al. 2009).

Winter severity and kernel density estimation

We estimated winter severity by summing daily snow pack estimates (measured in snow water equivalents [SWE], or the amount of water in a column of snow) from October 1 to April 30, based on the snow pack model described by Watson et al. (2009). We categorized winters during 1988-2008 as mild, average, or severe, with an average winter falling within the range of the median snowpack \pm 0.5 SD of the 30 years of data.

We performed spatial data manipulations and analyses using ArcGIS v9.3 (Environmental Systems Research Institute, Redlands, California). We obtained bison spatial information from aerial surveys conducted during 2000-2008 by the National Park Service. We grouped the data from these years by the previously-defined winter severity classifications to separate population spatial distributions for the three different types of winters. We then focused on the spatial locations during June when cattle were grazing on allotments (2000-2002, 2007-2008) and weighted spatial data points by the observed group size at that location.

We used Animal Space Use v1.3 (Horne and Garton 2009) to determine the appropriate bandwidth for our home range kernels. Next, we calculated home range distributions using a 95% fixed kernel estimator with Hawth's tools v3.27 (Rodgers and Carr 1998, Beyer 2004, Fieberg 2007). Because the primary season for cattle exposure began in June, we used the May-June spatial data.

We bootstrapped elk home-range kernels from minimum convex polygons representing the distributions of various elk populations (Hamlin and Cunningham 2008). We randomly assigned hypothetical individual animal locations within the bounds of the distribution. Then, subsequent points for each individual were approximated using spatial spread statistics on individual animal movements for elk from the northern Yellowstone herd (YNP, unpublished data). The bootstrapping was performed using R statistical language v2.11.0 (R Development Core Team 2010) and several R packages (Stabler 2006, Rowlingson et al. 2009, Lewin-Koh et al. 2010). Locations and usage of cattle grazing allotments were provided by the National Forest Service and the Animal and Plant Health Inspection Service. Our information on cattle grazing practices was limited to operations using public grazing allotments. We masked the home range

kernels to show specific overlap regions with cattle grazing allotments and calculated the percentage of volume overlap using R.

Statistical Analyses

We ran the risk model for 50,000 iterations to assess the variability of the exposure risk outputs. We determined the median number of abortion-days and infectious birth-days for both bison and elk. The numbers of abortion or infectious birth-days were defined as the number of infectious events (abortions or births) multiplied by the amount of days that each of those events will persist on the landscape. We determined the cattle exposure risk from both the Yellowstone bison herd as well as the five elk populations in the northern portion of the GYA. Probability intervals (95%) were estimated based on the 2.5th and 97.5th percentiles of the iterated values. Maps of the probability of infectious events across the northern portion of the GYA were made using ArcGIS. We also evaluated data from the Montana Department of Livestock on the number of bison migrating from YNP into Montana during 2000-2008 compared to the number of bison in YNP.

Model parameter sensitivity analysis

We performed a general sensitivity analysis of all input parameters in @RISK to determine which model parameters were most influential on *Brucella* exposure risk. In addition, the radius of exposure for an infectious event and wildlife tolerance factors were varied and the resultant change in the percentage of risk from bison exposure was evaluated. The model was updated for 50,000 iterations for each parameter value. Also, we evaluated the creation of bison home-range kernels by comparing the spatial distribution and associated risk of cattle *Brucella*

exposure between the kernel derived solely from June bison locations and a kernel derived from May and June locations.

RESULTS

Neither wildlife population had any projected infectious parturitions in January and February, so the only shedding in that time period was from abortions (Table 2). Bison began showing infectious parturitions in April, while all parturitions for elk were in May and June. Infectious event maps showed variable shedding across the northern GYA landscape (Fig. 2). Of the total number of infectious events in the northern Yellowstone elk population, 13.5% (95% P.I. = 2.5 to 45.0) were abortions as opposed to infectious parturitions. In bison, 16.5% (95% P.I. = 10.6 to 21.4) of the infectious events were abortions. The estimated annual cattle risk of exposure to a bison brucellosis infectious event was small (≤ 0.01 cattle-exposure event-days/year; Table 3). More risk was estimated in average and severe winters than for mild winters (Table 3). Two populations of elk in the northern portion of the GYA (Madison-Firehole and Sand Creek) had no detectable spatio-temporal overlap with cattle grazing allotments (see Supporting Information). Cattle risk estimates for exposure to an elk brucellosis infectious event were two orders of magnitude higher than for bison when elk range overlapped with cattle grazing allotments. Risk estimates for the Gallatin-Madison, Gravelly-Snowcrest, and Northern Yellowstone elk populations were 2.7, 1.6, and 1.9 cattle-exposure event-days, respectively. Bison migration data showed that the largest scale out-migration (2007-2008) occurred during a severe winter, with fewer animals in the YNP bison herd than the next largest migration (2005-2006) during an average winter (Table 4). Both of these migrations occurred during years when the Yellowstone bison herd was larger than 4400 animals.

Sensitivity analysis

The cattle exposure risk to a brucellosis event was most sensitive to the: (1) radius of exposure from each infectious event; (2) number of days infectious tissue would persist on the landscape prior to scavenging; (3) proportion of seropositive elk in the northern portion of the GYA; (4) proportion of elk shedding *Brucella* organisms; and (5) the adult female proportion of elk. Altering the radius of exposure by 50-meter increments from 50 to 250 meters yielded median exposure risks of 0.5 to 4.4 cattle-exposure event-days. Changing the wildlife tolerance factor from 0.1 to 0.9 caused the percentage of total risk attributable to bison to change from 0.1% to 0.6%. Using bison spatial locations from both May and June when deriving their home-range kernel increased greatly increased the amount of cattle exposure risk and increased the percentage of risk attributable to bison from $\leq 1\%$ using June alone to $\geq 60\%$.

DISCUSSION

Although our results support substantial shedding of *Brucella* bacteria from bison in some winters, the most substantial risk of bacterial transmission to cattle was from elk. Future risk estimates for bison depend on adaptive management of the population. Interactive effects between population size and winter severity were major determinants influencing bison movements to lower elevation winter grazing areas and overlap with federally-regulated domestic cattle grazing allotments. However, during the critical period of potential *B. abortus* exposure to cattle, the risk from Yellowstone bison was minimal. Natural movements of animals back to higher elevation summer ranges and boundary management operations were important in minimizing the contribution of bison to cattle exposure risk, which supports continued boundary

management operations for spatio-temporal separation between bison and cattle. Under current management practices, bison risk to cattle grazing in the northern portion of the GYA is expected to be small.

Maintaining spatial and temporal separation between bison and cattle, is believed to make the risk of *B. abortus* transmission from bison to cattle in the northern GYA negligible (Kilpatrick et al., 2009), but risk of transmission among bison remains high, accounting for the documented high prevalence (Cheville et al. 1998). Behavioral differences between species may also contribute to differences in pathogen prevalence. Spontaneous abortions by elk that are not segregated from their herd could expose many susceptible elk and cattle to infected fetuses and birth tissues (P.J. White, personal communication). In contrast, bison are gregarious during parturition, and pregnant females have been observed to nuzzle newborn calves (Treanor et al. 2008). Mobbing events of a newborn calf or aborted fetus could contribute to intra-species transmission of bacteria if the dam was infected (Jones et al. 2009).

Our results are consistent with the conclusion of Kilpatrick et al. (2009) that bison under current management practices are not likely to transmit *B. abortus* to cattle grazing in the northern portion of the GYA. However, we disagree with assuming that, under a “no plan” strategy (i.e., without management), the risk of bacterial transmission from bison to cattle would be low due to animal migration back to higher elevation grazing lands in Yellowstone National Park. This conclusion does not take into account the seasonal or environmental conditions, which may delay natural migration and does not consider that, without intensive management intervention, there is little doubt that bison would continue to expand their range and disperse to suitable habitat areas outside the northern and western boundaries of the park where cattle could come into contact with *Brucella* bacteria shed on birth tissues (Plumb et al. 2009). Lack of

consideration of boundary management operations makes accurate predictions of future spatial movements and locations of bison unlikely. The predictions and conclusions here are reasonable because bison are currently restricted to only a small fraction of their original range by active hazing into the Park as needed during the winter and spring to reduce contact with cattle. Spatial risk estimates are inextricably tied to current policy conditions and must be revisited as wildlife populations are adaptively managed.

The strength of our conclusions is based on the spatial and temporal resolution of the data used to parameterize the model. The cross-sectional nature of the bison aerial survey data limited our investigation to herd movement patterns. Also, the limited availability of appropriate elk location data, prohibited us from exploring how seasonal changes in elk distributions altered local risk of shedding.

This model provides the first spatially-explicit framework for assessing the risk of bacterial shedding of *B. abortus* by bison and elk across the northern portion of the GYA. It may be expanded to include the entire GYA, or serve as a template for models of other diseases. The next steps in exploring the risk of *B. abortus* transmission in the northern GYA are to continue to refine our model with new data, especially on spatial locations of cattle and wildlife, as well as animal movements. The underlying disease dynamics between elk and bison also need to be examined to estimate what frequency or rate of interspecies pathogen transmission is necessary to be maintaining the current prevalence in elk and bison populations in the northern GYA and relative impact that alternative management strategies can have on overall transmission.

In addition to overlap, the major contributors to risk were wildlife population size and the number of elk that were shedding *Brucella* bacteria. While elk currently have a lower density of shedding events throughout their range, they have a larger spatio-temporal overlap with cattle

and are more tolerated by managers and livestock keepers on public grazing allotments. Thus, the predominant source of risk to cattle in the northern portion of the greater Yellowstone area is from elk. With increased disease prevalence due to increased winter densities or other factors, elk are likely to contribute greatly to the overall level of bacterial shedding on the northern GYA landscape (Fig. 2) and will continue to represent the vast majority of risk of *B. abortus* exposure to cattle grazing in the northern portion of the GYA. Therefore, brucellosis management efforts should focus more on the comingling of cattle and elk during the critical abortion period to more effectively decrease risk of transmission.

ACKNOWLEDGEMENTS

Funding for this project was provided by the US Department of Agriculture: Animal and Plant Health Inspection Service (APHIS) and the US National Park Service through the Yellowstone Wildlife Health Program. The project also received support from the Foreign Animal and Zoonotic Disease Defense Center of Excellence, through a grant from the Department of Homeland Security, Science and Technology Directorate, Office of University Programs. We thank all collaborators who provided data for parameterization of the risk model, including Ken Britton and Lisa Stoeffler from National Forest Service, Mark Drew from the Idaho Department of Fish and Game, and Rebecca Frey from APHIS-Veterinary Service. We also thank Shane Grube from the Montana Department of Livestock for compiling data on bison in Montana; Keith Aune from the Wildlife Conservation Society and Brandon Scurlock and Hank Edwards from the Wyoming Game and Fish Department for suggestions on model improvement; Lindsey Holmstrom from the Center for Animal Disease Modeling and Surveillance (CADMS) for help

with GIS analysis; and Glenn Plumb and P.J. White for their intellectual contributions and for pursuing funding and building the collaborative relationships that made this work possible.

REFERENCES

- Aune, K., J. C. Rhyan, B. Corso, and T. Roffe. 2007. Environmental persistence of *Brucella* organisms in natural environments of the greater Yellowstone area - A preliminary analysis. Report of the Committee on Brucellosis. United States Animal Health Association, Richmond, Virginia.
- Barber-Meyer, S. M., P. J. White, and L. D. Mech. 2007. Survey of selected pathogens and blood parameters of northern Yellowstone elk: wolf sanitation effect implications. *American Midland Naturalist* **158**:369-381.
- Berger, J., and S. L. Cain. 1999. Reproductive synchrony in brucellosis-exposed bison in the southern greater Yellowstone ecosystem and in noninfected populations. *Conservation Biology* **13**:357-366.
- Beyer, H. L. 2004. Hawth's Analysis Tools for ArcGIS.
<<http://www.spataleecology.com/htools>>.
- Cheville, N., D. R. McCullough, and L. R. Paulson. 1998. Brucellosis in the greater Yellowstone area. National Research Council, Washington D.C.
- Clarke, R., C. Jourdonnais, J. Munding, L. Stoeffler, and R. Wallen. 2005. A Status Review of Adaptive Management Elements 2000 to 2005. Interagency Bison Management Plan.
- Coe, P. K., B. K. Johnson, K. M. Stewart, and J. G. Kie. 2005. Spatial and temporal interactions of elk, mule deer and cattle. Pages 150-158 in M. J. Wisdom, editor. *The Starkey Project: a synthesis of long-term studies of elk and mule deer*. Reprinted from the 2004 Transactions of the North American Wildlife and Natural Resources Conference. Alliance Communications Group, Lawrence, Kansas, USA.

- Cook, W. E., E. S. Williams, and S. A. Dubay. 2004. Disappearance of bovine fetuses in northwestern Wyoming. *Wildlife Society Bulletin* **32**:254-259.
- Cross, P. C., T. O. Lemke, P. J. White, and D. B. Tyers. 2009. Northern Yellowstone cooperative wildlife working group 2008 annual report (October 1, 2007-September 30, 2008). U.S. Geological Survey, Northern Rocky Mountain Science Center, Bozeman, Montana.
- Davis, D., J. W. Templeton, T. A. Ficht, J. D. Williams, J. D. Kopec, and L. G. Adams. 1990. *Brucella abortus* in captive bison. I. Serology, bacteriology, pathogenesis, and transmission to cattle. *Journal of Wildlife Diseases* **26**:360-371.
- Delahay, R. J., G. C. Smith, and M. R. Hutchings. 2009. The Science of Wildlife Disease Management. Pages 1-8 in R. J. Delahay, G. C. Smith, and M. R. Hutchings, editors. *Management of Disease in Wild Mammals*. Springer, Tokyo, Japan.
- Dobson, A. 2004. Population Dynamics of Pathogens with Multiple Host Species. *The American Naturalist* **164**:S64-S78.
- Donch, D. A., and A. A. Gertonson. 2008. Status report -- Fiscal year 2008; Cooperative State-Federal Brucellosis Eradication Program. USDA-APHIS Veterinary Services.
- Donch, D. A., A. A. Gertonson, and M. J. Gilsdorf. 2005. Cooperative State-Federal Brucellosis Eradication Program-Status Report-Fiscal Year 2004. The Annual Meeting of the United States Animal Health Association. United States Animal Health Association.
- Eberhardt, L. L., P. J. White, R. A. Garrott, and D. B. Houston. 2007. A seventy-year history of trends in Yellowstone's northern elk herd. *Journal of Wildlife Management* **71**:594-602.
- Fieberg, J. 2007. Utilization distribution estimation using weighted kernel density estimators. *Journal of Wildlife Management* **71**:1669-1675.

- Godfroid, J. 2002. Brucellosis in wildlife. *Revue Scientifique et Technique Office International des Epizooties* **21**:277-286.
- Government Accountability Office. 2009. Veterinarian Workforce: Actions are Needed to Ensure Sufficient Capacity for Protecting Public and Animal Health. Government Accountability Office, Washington, D.C.
- Hamlin, K. L. 2006. Monitoring and Assessment of Wolf-Ungulate Interactions and Population Trends within the Greater Yellowstone Area, Southwestern Montana, and Montana Statewide. *Montana Fish, Wildlife & Parks*.
- Hamlin, K. L., and J. A. Cunningham. 2008. Montana elk movements, distribution, and numbers relative to brucellosis transmission risk. *Montana Fish, Wildlife, and Parks*.
- Horne, J. S., and E. O. Garton. 2009. Animal Space Use 1.3.
<http://www.cnr.uidaho.edu/population_ecology/animal_space_use>.
- Johnson, D. E. 1951. Biology of the elk calf, *Cervus canadensis nelsoni*. *Journal of Wildlife Management* **15**:396-410.
- Jones, J. D., J. J. Treanor, and R. L. Wallen. 2009. Parturition in Yellowstone bison. National Park Service, Yellowstone National Park, Wyoming.
- Kilpatrick, A. M., C. M. Gillin, and P. Daszak. 2009. Wildlife-livestock conflict: the risk of pathogen transmission from bison to cattle outside Yellowstone National Park. *Journal of Applied Ecology* **46**:476-485.
- Kittelberger, R., F. Hilbank, M. F. Hansen, M. Penrose, G. W. de Lisle, J. J. Letesson, B. Garin-Bastuji, J. Searson, C. A. Fossati, A. Cloeckert, and G. G. Schurig. 1995. Serological crossreactivity between *Brucella abortus* and *Yersinia enterocolitica* O:9 I immunoblot

- analysis of the antibody response to *Brucella* protein antigens in bovine brucellosis. *Veterinary Microbiology* **47**:257-270.
- Lewin-Koh, N. J., R. Bivand, contributions by Edzer J. Pebesma, E. Archer, A. Baddeley, H.-J. r. Bibiko, S. p. Dray, D. Forrest, M. Friendly, P. Giraudoux, D. Golicher, V. G. m. Rubio, P. Hausmann, T. Jagger, S. Luque, D. MacQueen, A. Niccolai, T. Short, and B. Stabler. 2010. maptools: Tools for reading and handling spatial objects. R package version 0.7-34.
- Maichak, E. J., B. M. Scurlock, J. D. Rogerson, L. L. Meadows, A. E. Barbknecht, W. H. Edwards, and P. C. Cross. 2009. Effects of management, behavior, and scavenging on risk of brucellosis transmission in elk of western Wyoming. *Journal of Wildlife Diseases* **45**:398-410.
- Meagher, M. 1973. The bison of Yellowstone National Park. National Park Service, Washington, D.C., USA.
- Meagher, M., and M. E. Meyer. 1994. On the origin of brucellosis in bison of Yellowstone National Park: A review. *Conservation Biology* **8**:645-653.
- Mohler. 1917. Report of the Chief of the Bureau of Animal Industry, Pathological Division. United States Department of Agriculture, Washington, D.C.
- Peterson, M. J., W. E. Grant, and D. S. Davis. 1991. Bison-Brucellosis Management - Simulation of Alternative Strategies. *Journal of Wildlife Management* **55**:205-213.
- Plumb, G. E., P. J. White, M. B. Coughenour, and R. L. Wallen. 2009. Carrying capacity, migration, and dispersal in Yellowstone bison. *Biological Conservation* **142**:2377-2387.
- Power, A. G., and C. E. Mitchell. 2004. Pathogen Spillover in Disease Epidemics. *The American Naturalist* **164**:S79-S89.

- R Development Core Team. 2010. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Rodgers, A. R., and A. P. Carr. 1998. HRE: The Home Range Extension for ArcView. Ontario Ministry of Natural Resources, Centre for Northern Forest Ecosystem Research, Thunder Bay, Ontario, Canada.
- Roffe, T. J., J. C. Rhyon, K. Aune, L. M. Philo, D. R. Ewalt, T. Gidlewski, and S. G. Hennager. 1999. Brucellosis in Yellowstone National Park bison: Quantitative serology and infection. *Journal of Wildlife Management* **63**:1132-1137.
- Rowlingson, B., P. Diggle, adapted, packaged for R by Roger Bivand, pcp functions by Giovanni Petris, and goodness of fit by Stephen Eglen. 2009. splancs: Spatial and Space-Time Point Pattern Analysis. R package version 2.01-25.
- Stabler, B. 2006. shapefiles: Read and Write ESRI Shapefiles. R package version 0.6.
- Thorne, E. T., J. K. Morton, F. M. Blunt, and H. A. Dawson. 1978. Brucellosis in elk. II. Clinical effects and means of transmission as determined through artificial infections. *Journal of Wildlife Diseases* **14**:280-291.
- Treanor, J., J. Johnson, R. Wallen, S. Cilles, P. Crowley, and D. Maehr. 2008. Vaccination strategies for managing brucellosis in Yellowstone bison. National Park Service, Yellowstone National Park, Wyoming.
- U.S. Department of Interior [USDI], and U.S. Department of Agriculture [USDA]. 2000. Record of Decision for Final Environmental Impact Statement and Bison Management Plan for the State of Montana and Yellowstone National Park. USDI National Park Service and USDA Animal and Plant Health Inspection Service.

Watson, F. G. R., T. N. Anderson, W. B. Newman, S. S. Cornish, and T. R. Thein. 2009.

Modeling Spatial Snow Pack Dynamics. Pages 85-112 *in* R. A. Garrot, P. J. White, and

F. G. R. Watson, editors. *The Ecology of Large Mammals in Central Yellowstone:*

Sixteen Years of Integrated Field Studies. Elsevier, New York.

Woolhouse, M. E. J., L. H. Tayloer, and D. T. Haydon. 2001. Population Biology of Multihost

Pathogens. *Science* **292**:1109-1112.

Table 1. Input parameters for a *Brucella abortus* transmission model used to assess the risk of an infectious event occurring in elk and bison populations in the northern greater Yellowstone area.

Description of variables	Statistical distribution (parameters) [Mean, SD]	Source
Shedding proportion	Beta (12,14) [0.46, 0.10]	(Roffe et al. 1999) ^a
Fetal persistence	BetaGeneral (2, 6.93, 1, 78) [18.25, 10.19]	(Aune et al. 2007)
<u>Bison</u>		
Number of animals Fit from 2000-2008 data	Logistic (3788.53, 450.13) [3788.53, 816.45]	(National Park Service, unpublished data)
Age proportion (of total population): Fit from 2004-2008 data		(National Park Service, unpublished data)
2-3 year-old females	BetaSubjective (0.043, 0.047, 0.04736, 0.053) [0.047, 0.002]	
4+ year-old females	Pareto (46.43, 0.35123) [0.36, 0.01]	
Proportion pregnant:		
2-3 year-old	Uniform (0.71, 0.79) [0.75, 0.02]	(Yellowstone Center for Resources 2008)
4+ year-old	Uniform (0.76, 0.89) [0.83, 0.04]	
Proportion seropositive 2+ year-old <u>(sampled at boundary capture facility)</u>	Beta (331.0, 211.6) [0.61, 0.02]	(National Park Service, unpublished data)

Percentage shedding by abortion:

First pregnancy females	BetaSubjective (0.65, 0.78, 0.78, 0.9) [0.78, 0.07]	(Davis et al. 1990)
Mature females	BetaSubjective (0.01, 0.1, 0.09, 0.15) [0.09, 0.03]	(Peterson et al. 1991)
Birth synchrony	Normal (40.57, 13.33) [40.57, 13.33] Day 1 = April 1	(Berger and Cain 1999)

Elk

Adult female proportion Fit from 2000-2008 data	BetaSubjective (0.52, 0.73, 0.7, 0.8) [0.7, 0.06]	(National Park Service, unpublished data)
Adult female: yearling	10:1	(National Park Service, unpublished data)
Proportion pregnant: Fit from 2000-2006 data		(National Park Service, unpublished data)
Yearling	BetaSubjective (0.1, 0.33, 0.32, 0.4) [0.32, 0.03]	
Adult	BetaSubjective (0.78, 0.82, 0.815, 0.84) [0.82, 0.01]	
Percentage of shedding by abortion:		
First pregnancy females	Beta (13.3, 14.4) [0.48, 0.09]	(Thorne et al. 1978)
Mature females	Beta (1.2, 6.8) [0.15, 0.12]	
Birth synchrony	Poisson (32.526) [32.526, 5.703] Day 1 = May 1	(Maichak et al. 2009)

Gallatin-Madison

Number of animals	Normal (7807, 793)	
Fit from 2000-2008 estimates (sightability corrected using 1.322 correction factor)	[7807, 793]	(Hamlin and Cunningham 2008)

B. abortus seropos. proportion Beta (3.1, 101.5)
[0.03, 0.02]

Gravelly-Snowcrest

(Hamlin 2006)

Number of animals Uniform (10,900, 11,570)
Fit from 2004&2006 data [11,235, 193]

B. abortus seropos. proportion Beta (3.1, 101.5)
[0.03, 0.02]

Madison-Firehole

Number of animals Loglogistic (236.8, 196.2, 1.4)
Fit from 2000-2008 estimates [757.2, N/A] (Hamlin and Cunningham 2008)
(sightability corrected using 1.322 correction factor)

B. abortus seropos. proportion Beta (3.1, 101.5)
[0.03, 0.02]

Northern Yellowstone

Number of animals Lognormal (9742, 3801, Shift (3396))
Fit from 2000-2008 estimates [13,137, 3800] (Cross et al. 2009)
(sightability corrected using 1.322 correction factor)

B. abortus seropos. proportion Uniform (0.01, 0.05) (Barber-Meyer et al. 2007)
[0.03, 0.01]

Sand Creek, Idaho

Number of adult females, 2006 1,413 (Mark Drew, Idaho
(sightability corrected using 1.322 correction factor) Department of Fish and
Game, unpublished data)

B. abortus seropos. proportion Beta (0.9, 100)
[0.01, 0.01]

a – study generalizes statistic for seropositive female bison

Table 2. Median number of abortion-days and infectious parturition-days and 95% probability intervals (P.I.) for bison and elk in the northern portion of the greater Yellowstone area.

Season	Bison		Elk	
	Abortion-days (95% P.I.)	Infectious Parturition-days (95% P.I.)	Abortion-days (95% P.I.)	Infectious Parturition-days (95% P.I.)
Jan.-Feb.	3.3 (0.6, 10.9)	0 (0, 0)	0.01 (0.00, 0.06)	0 (0, 0)
Mar.-Apr.	9.4 (1.7, 31.2)	16.9 (3.2, 52.7)	4.7 (0.4, 37.7)	0 (0, 0)
May-Jun.	2.2 (0.3, 6.1)	58.7 (11.0, 182.7)	2.9 (0.2, 23.2)	50.1 (6.5, 239.8)
Total	14.9; (2.6, 48.2)	75.6 (14.2, 235.4)	7.6 (0.6, 61.0)	50.1 (6.5, 239.8)
% of total shedding	16.5% (10.6, 21.4)	83.5% (78.6, 89.4)	13.5% (2.5, 45.0)	86.5% (55.0, 97.5)

Table 3. Median cattle risk of exposure to a *Brucella abortus* infectious shedding event from the Yellowstone bison population for the month of June using home range estimates for mild, average, and severe winters with 95% probability intervals (P.I.). The units for risk are cattle-exposure event-days.

	Cattle Risk of <i>Brucella</i>	% of Total Exposure Risk
	Transmission	From Bison
	(95% P.I.)	(95% P.I.)
Mild	<0.01 (0, 0.01)	<0.1 (0, 0.2)
Average	0.01 (0, 0.12)	0.3 (0, 1.8)
Severe	0.01 (0, 0.13)	0.3 (0, 2.1)

Table 4. Yellowstone bison population estimates and the corresponding number of out-migrating bison in the Western Management Area (WMA) during the month of February for mild (M), average (A), and severe (S) winters from 1999 to 2008.

Year	Bison	Bison in WMA (February)
1999-2000 (A)	2500	1
2000-2001 (M)	3000	10
2001-2002 (A)	3400	6
2002-2003 (A)	4100	10
2003-2004 (A)	4250	1
2004-2005 (M)	4400	2
2005-2006 (A)	5000	157
2006-2007 (M)	4000	No Data
2007-2008 (S)	4700	182

Fig. 1. Probability distributions for infectious parturitions and abortions by bison and elk in the northern portion of the greater Yellowstone area.





Fig. 2. Map of total *Brucella abortus* shedding events during June in the northern portion of the greater Yellowstone area based on an average winter. Red areas indicate higher levels of shedding while yellow areas indicate lower levels of shedding.



Online Supporting Information

Table S1. Median cattle risk of exposure to a *Brucella abortus* infectious shedding event from five elk populations in the northern portion of the greater Yellowstone area and 95% probability intervals (P.I.). The units for this risk are given as cattle-exposure event-days/year.

Elk Population	Cattle Risk of <i>Brucella</i> Exposure (95% P.I.)
Gallatin-Madison	2.7 (0.1, 27.0)
Gravelly-Snowcrest	1.6 (0.1, 15.8)
Madison-Firehole	0.0 (0.0, 0.0)
Northern Yellowstone	1.9 (0.1, 20.3)
Sand Creek	0.0 (0.0, 0.0)

Fig. S1. Risk model equations for: (a) risk of cattle exposure to a wildlife brucellosis infectious event; (b) total infectious event-days, (b) abortion-days, and (c) infectious live parturition-days from elk and bison in the northern portion of the greater Yellowstone area.

(a)

Risk \propto

$$\begin{aligned} & (\text{Number of cattle on allotment}) \times (\text{Number of days cattle are at-risk}) \times \\ & (\text{Number of wildlife infectious event-days}) \times [(\text{Area of Event Exposure}) / (\text{Area of} \\ & \text{Allotment})] \times (\text{Wildlife Tolerance Factor}^\dagger) \end{aligned}$$

(b)

Number of infectious event-days =

$$(\text{Number of abortion-days}) + (\text{Number of infectious live parturition-days})$$

(c)

Number of abortion-days =

$$\begin{aligned} & (\text{Number of animals}) \times \\ & [(\text{Proportion first pregnancy}) \times (\text{Age-specific pregnancy proportion}) \\ & \times (\text{Age-specific shedding proportion}) \times (\text{proportion of first pregnancy females aborting}) + \\ & (\text{Proportion mature females}) \times (\text{Age-specific pregnancy proportion}) \\ & \times (\text{Age-specific shedding proportion}) \times (\text{proportion of mature females aborting})] \\ & \times (\text{Proportion of total abortions expected to occur in the time window}) \end{aligned}$$

x (Bacterial persistence proportion)

(d)

Number of infectious live parturition-days =

(Number of animals) x

[(Proportion first pregnancy) x (Age-specific pregnancy proportion)

x (Age-specific shedding proportion) x (proportion of first pregnancy females not aborting)]

+

[(Proportion mature females) x (Age-specific pregnancy proportion)

x (Age-specific shedding proportion) x (proportion of mature females not aborting)]

x (Proportion of total infectious live-parturitions expected to occur in the time window)

x (Bacterial persistence proportion)

† - Defined as the access that bison have on grazing allotments as a percentage of the access that elk are given to allotments.

Objective 2

Who infects whom? Interspecies transmission dynamics of brucellosis in the northern greater Yellowstone area

Brant A. Schumaker,^{1*} Jonna A.K. Mazet,² John Treanor,³ Rick Wallen,³ Ian A. Gardner,⁴ Martin Zaluski,⁵ and Tim E Carpenter¹

¹Center for Animal Disease Modeling and Surveillance (CADMS), Department of Medicine and Epidemiology, School of Veterinary Medicine, University of California, Davis, CA 95616, USA;

²Wildlife Health Center, School of Veterinary Medicine, University of California, Davis, CA 95616, USA; ³National Park Service, Yellowstone National Park, P.O. Box 168, Wyoming

82190, USA; ⁴Department of Medicine and Epidemiology, School of Veterinary Medicine,

University of California, Davis, CA 95616, USA; ⁵Montana Department of Livestock, P.O. Box 202001, Helena, MT 59620-2001, USA;

*Corresponding author (email: theschu@ucdavis.edu, phone: (530) 752-3566, fax: (530) 752-1618)

Abstract

Bison (*Bison bison*) and elk (*Cervus elaphus*) in the northern portion of the greater Yellowstone area (GYA) remain reservoirs capable of transmitting *Brucella abortus* bacteria to livestock. However, the inter- and intra-species contact rates required to maintain brucellosis in the GYA have not previously been characterized. Without this knowledge, the likely effects of risk mitigation strategies cannot be adequately evaluated. We used a risk model to estimate the spatio-temporal distribution of *B. abortus* shedding events from bison and elk populations in the northern GYA. The percentage of *B. abortus* infectious events in overlapping wildlife populations was calculated, and the risk of *B. abortus* transmission within and between populations was estimated. Bison risk from other bison and from elk showed almost 100% adequacy to transmit the organism once spatio-temporal overlap occurred; however, contact within elk populations was only approximately 34% adequate. Transmission risks to elk from elk in other populations or from bison were very small. Minimal opportunity exists for *B. abortus* transmission from bison to elk under current natural conditions in the northern GYA. Under current conditions, management alternatives that reduce bison seroprevalence are unlikely to substantially reduce transmission risk from elk to cattle. Strategies that decrease elk herd densities and group sizes and reduce elk-to-elk transmission could reduce the overall risk to cattle grazing in the northern portion of the GYA.

Keywords: bison, Brucella abortus, infectious disease, model, elk, population management, risk, transmission, wildlife

INTRODUCTION

Bison (*Bison bison*) and elk (*Cervus elaphus*) populations in the northern greater Yellowstone area (GYA) are variably infected with *Brucella abortus*. Elk populations in the northern GYA have a relatively low seroprevalence (i.e., exposure; <5%) of *B. abortus*, whereas seroprevalence in Yellowstone bison is high (40-60%) (Hobbs et al. 2009). While bison most likely acquired brucellosis from cattle grazing in the GYA (Meagher and Meyer 1994), *B. abortus* has been eradicated from livestock in the US, and wildlife in the northern greater Yellowstone area (GYA) remain a source for potentially transmitting *B. abortus* bacteria to livestock.

The Interagency Brucellosis Management Plan (IBMP) was established in 2000 to manage the risk of *B. abortus* transmission from bison to cattle by implementing hazing, test-and-slaughter, hunting, and other actions near the boundary of Yellowstone National Park (YNP) (Plumb and Aune 2002). To date, these actions have successfully prevented the transmission of *B. abortus* from bison to cattle (Clarke et al. 2005), and assessments suggest the risk of future *B. abortus* transmission is minimal under current management conditions (Kilpatrick et al. 2009, Schumaker et al. 2010). Conversely, elk in the northern GYA have received relatively little brucellosis management attention until recently and often move freely across the ecosystem and come into close contact with cattle. All detections of *B. abortus* infection in northern GYA cattle in the last decade have been attributed to elk (Donch and Gertonson 2008).

Having multiple hosts increases the complexity of *B. abortus* transmission dynamics (Dobson 2004, Delahay et al. 2009). There is still an insufficient understanding of much of these dynamics, and this information is crucial for disease management. Elk with significant home-range overlap with Yellowstone bison do not show evidence of an increase in *B. abortus*

exposure compared with populations with spatio-temporal separation from bison (Ferrari and Garrot 2002, Proffitt et al. 2010). Apparently lower *B. abortus* exposure in elk may be due to differences in the immunological responses or reproductive behavior of the wildlife hosts. Without better knowledge of the inter- and intra-species contact rates that maintain *B. abortus* prevalence in the GYA, the likely effects of risk mitigation strategies cannot be evaluated thoroughly.

The Yellowstone bison population has been extensively modeled (Peterson et al. 1991, Dobson and Meagher 1996, Gross et al. 2002, Treanor et al. 2010). However, none of these models attempt to estimate the contact rates required to maintain *B. abortus* at documented prevalence levels. We extended past modeling efforts by quantifying the transmission dynamics within and between elk and bison populations in the northern GYA. We also determined the bison and elk intra- and inter-species contact rates required to maintain documented prevalence levels in elk in the northern GYA.

MATERIALS AND METHODS

Study area and wildlife host populations

The GYA is one of the largest intact temperate zone ecosystems on earth and also home to the largest wild and free-ranging elk and bison populations in the United States. One bison population with between 2,000 and 5,000 individuals (Meagher 1973, Clarke et al. 2005) and five elk populations – Gallatin-Madison (GM), Gravelly-Snowcrest (GS), Madison-Firehole (MF), northern Yellowstone (NY), and Sand Creek, Idaho (SC) are distributed across 3,000 km² in the northern GYA. Estimates of northern Yellowstone elk were near 25,000 animals in the late 1980s, but decreased by approximately 50-60% by 2006 (Eberhardt et al. 2007). Median

estimates fit from multiple data sets for the other four elk populations were: 7,807 for GM; 11,253 for GS; 757 for MF; and 1,413 for SC, respectively (Table 1).

Risk model

The estimation of the *B. abortus* transmission potential within and between elk and bison populations in the northern GYA employed a previously developed risk model (Schumaker et al. 2010). The model estimated the number and spatiotemporal distribution of *B. abortus* shedding events from third-trimester abortions and infectious live parturition events from one bison and five elk populations in the northern GYA (Figure 1). The stochastic model was parameterized with statistical distributions fit to winter severity, animal location, serologic testing, demographic and epidemiologic data using @RISK v5.5 (Palisade Corporation, Ithaca, New York, USA; Table 1). The assumptions for the model were: 1) adult females are the primary source of infection; 2) the critical season of transmission is between January 1 and June 30; 3) no fully immune state exists; and 4) random mixing of animals occurs within a population.

Risk calculation

Risk of *B. abortus* transmission is a combination of spatiotemporal overlap of at-risk individuals, the number and location of infectious events within their own or a neighboring population of bison or elk, and behavioral and disease factors that allow transmission within and between wildlife populations (Equation 1). These factors could include the relative dominance of one species over another, which might include driving a group of animals off grazing land where infectious material could reside. They also could include transmission rates for *B. abortus* within and between elk and bison, respectively, once exposed to the pathogen.

Numbers of infectious events in each wildlife population were taken from the results of 50,000 iterations of the stochastic risk model and distributed to fixed kernel density estimations of wildlife home ranges as described elsewhere (Schumaker et al. 2010). Overlap among elk and bison populations was calculated using the Spatial Analyst extension in ArcGIS v9.3 (Environmental Systems Research Institute, Redlands, California). Rasters were converted to ASCII files and the percentage of volume overlap was calculated using R statistical language v2.11.1 (R Development Core Team 2010) and the raster R package (Hijmans and Van Etten 2010).

Statistical analyses

Equations were created for *B. abortus* transmission risk using medians of the risk model parameters (Table 2). The probabilities of adequate contact – contact, which would result in transmission if the exposed animal were susceptible to infection – from spatiotemporal overlaps of wildlife with *B. abortus* infectious events were listed as unknowns. These probabilities were estimated from the data as uniform distributions, using the observed minimum and maximum values. The Solver Add-in for Excel v2007 (Microsoft Inc., Redlands, WA, USA) was used to optimize a solution using 0.000001 precision, 5% tolerance, 0.0001 convergence, tangent estimates, forward derivatives, and Newton search algorithm. Then the risk model was run for 50,000 iterations to assess the distribution of transmission risk within and between GYA wildlife populations. The median was determined and 95% probability intervals were estimated based on the 2.5th and 97.5th percentiles of the iterated values. *B. abortus* incidence in the wildlife populations was also estimated (Equation 2).

RESULTS

Bison overlaps within their own population and with elk were almost 100% adequate for *B. abortus* transmission, while elk overlaps with bison shedding were less than 0.1% adequate. Elk overlaps within their own population ranged from 33.8-34.0% adequate for *B. abortus* transmission. Elk overlaps with elk from other populations were only 1.4-1.6% adequate for *B. abortus* transmission, but 24 to 60 times more adequate for transmission than potential contacts from bison. As a percentage of total risk, bison transmission risk from within their own population was three times higher than from elk (Table 3). Conversely, elk risk from bison ranged from <0.1 – 0.5% of total risk. In the GM, GS, and NY populations, risk from other elk populations ranged from 0.3-7.7% of total risk. In those populations, risk from within their own population ranged from 92.1-99.4% of total risk. However, in the MF and SC populations risk from within their own population was only 27.1 and 39.9%, respectively, compared to 72.3 and 59.9% from other elk populations.

DISCUSSION

This study found that minimal opportunity exists for *B. abortus* transmission from bison to elk under natural conditions in the northern GYA. The reasons for this lower probability of adequate contact for *B. abortus* transmission, even when spatiotemporal overlap occurred, are likely immunological or behavioral. Differences in the immune systems of elk compared with bison may make them less susceptible to infection. These immunological differences may also account for the different responses of elk and bison to vaccination, leading to the failure of elk to be protected by RB51 vaccination while bison acquire some protection from the vaccine (Kreeger et al. 2002, Olsen et al. 2003). Also, anecdotally, bison are more dominant than elk and

may drive elk off grazing areas, increasing their opportunity for exposure to elk infectious material but decreasing the opportunity for elk to be exposed to bison infectious material (Rick Wallen, personal communication).

In addition, reproductive behavioral differences likely account for decreased transmission risk for elk compared with bison. The probability of *B. abortus* transmission between elk (or from elk to cattle) is likely low during calving (May through June) because pregnant dams isolate themselves while giving birth and meticulously clean the birth site (Johnson 1951). Thus, birth sites are dispersed, and the likelihood of other elk encountering infected birth tissues and fluids is low. However, transmission risk may be higher during the potential abortion period from February through April when many elk aggregate in larger groups on lower-elevation winter ranges that sometimes include ranch areas with cattle (Hamlin and Cunningham 2008). Spontaneous abortions by elk that are not segregated from their herd could expose many elk to infected fetuses and birth tissues (P.J. White, personal communication). In contrast, bison are gregarious during parturition, and pregnant females have been observed to nuzzle newborn calves (Yellowstone Center for Resources 2008). Mobbing events of a newborn calf or aborted fetus could contribute to intra-species transmission of bacteria if the dam were infected (Yellowstone Center for Resources 2009).

The MF and SC elk populations had the lowest estimated transmission risk. The SC population was spatio-temporally distant from the YNP bison herd, while the MF elk had increased overlap with Yellowstone bison (Ferrari and Garrot 2002, Proffitt et al. 2010). A lower median seroprevalence in the SC population (0.01 compared with 0.03 in all other elk populations) and the small population size in both herds resulted in decreased estimated shedding

in these populations. Therefore, a higher percentage of total risk to these populations came from outside elk sources rather than in the other three elk populations.

Probabilities for elk having adequate contact with other elk for *B. abortus* transmission were 24 times higher within their own population than from other elk populations. Because, behaviorally, most risk comes from spontaneous abortions, it is understandable that these abortions occur more frequently within a single elk population than during periods of comingling of multiple populations. For bison, transmission risk could potentially come from within their own population or from GYA elk. However, because there was a single population of bison, it decreased the ability to differentiate the relative probabilities for adequate contact from bison or elk shedding.

Estimates for transmission risk and transmission incidence were of the same order of magnitude. However, the equation for incidence used the lifespan of the wildlife species as the duration of infection. The assumption that all infected animals were infected at birth created an overestimate of duration, which resulted in an underestimate of incidence. This helps to account for a transmission incidence lower than the total transmission risk in bison as well as two populations of elk.

The National Park Service is exploring the remote delivery of the RB51 brucellosis vaccine to female Yellowstone bison to reduce abortions from this non-native disease and increase tolerance for bison outside YNP (USDI-NPS 2010). Vaccination is expected to significantly reduce the population seroprevalence of *B. abortus* infection (Yellowstone Center for Resources 2008). However, because bison rarely transmit *B. abortus* to elk, management alternatives such as vaccination that reduce bison seroprevalence are unlikely to reduce transmission from elk to cattle. However, these practices would increase the tolerance for bison

outside YNP boundaries, as they would decrease the potential for transmission from bison to cattle. The reduction in practices that increase elk herd densities and group sizes or the implementation of strategies to reduce elk-to-elk transmission should be promoted to reduce the overall risk to cattle grazing in the northern GYA.

ACKNOWLEDGEMENTS

Funding for this project was provided by the US Department of Agriculture: Animal and Plant Health Inspection Service (APHIS) and the US National Park Service through the Yellowstone Wildlife Health Program. The project was also funded through the Foreign Animal and Zoonotic Disease Defense Center of Excellence, by a grant from the Department of Homeland Security, Science and Technology Directorate, Office of University Programs. We thank Dr. Glenn Plumb and Dr. P.J. White for pursuing funding and building the collaborative relationships that made this work possible.

LITERATURE CITED

- Aune, K., J. C. Rhyan, B. Corso, and T. Roffe. 2007. Environmental persistence of *Brucella* organisms in natural environments of the greater Yellowstone area - A preliminary analysis. Report of the Committee on Brucellosis. United States Animal Health Association, Richmond, Virginia.
- Barber-Meyer, S. M., P. J. White, and L. D. Mech. 2007. Survey of selected pathogens and blood parameters of northern Yellowstone elk: wolf sanitation effect implications. *American Midland Naturalist* **158**:369-381.
- Berger, J., and S. L. Cain. 1999. Reproductive synchrony in brucellosis-exposed bison in the southern greater Yellowstone ecosystem and in noninfected populations. *Conservation Biology* **13**:357-366.
- Clarke, R., C. Jourdonnais, J. Munding, L. Stoeffler, and R. Wallen. 2005. A Status Review of Adaptive Management Elements 2000 to 2005. Interagency Bison Management Plan.
- Cross, P. C., T. O. Lemke, P. J. White, and D. B. Tyers. 2009. Northern Yellowstone cooperative wildlife working group 2008 annual report (October 1, 2007-September 30, 2008). U.S. Geological Survey, Northern Rocky Mountain Science Center, Bozeman, Montana.
- Davis, D., J. W. Templeton, T. A. Ficht, J. D. Williams, J. D. Kopec, and L. G. Adams. 1990. *Brucella abortus* in captive bison. I. Serology, bacteriology, pathogenesis, and transmission to cattle. *Journal of Wildlife Diseases* **26**:360-371.
- Delahay, R. J., G. C. Smith, and M. R. Hutchings. 2009. The Science of Wildlife Disease Management. Pages 1-8 in R. J. Delahay, G. C. Smith, and M. R. Hutchings, editors. *Management of Disease in Wild Mammals*. Springer, Tokyo, Japan.

- Dobson, A. 2004. Population Dynamics of Pathogens with Multiple Host Species. *The American Naturalist* **164**:S64-S78.
- Dobson, A., and M. Meagher. 1996. The Population Dynamics of Brucellosis in the Yellowstone National Park. *Ecology* **77**:1026-1036.
- Donch, D. A., and A. A. Gertonson. 2008. Satus report -- Fiscal year 2008; Cooperative State-Federal Brucellosis Eradication Program. USDA-APHIS Veterinary Services.
- Eberhardt, L. L., P. J. White, R. A. Garrott, and D. B. Houston. 2007. A seventy-year history of trends in Yellowstone's northern elk herd. *Journal of Wildlife Management* **71**:594-602.
- Ferrari, M. J., and R. A. Garrot. 2002. Bison and Elk: Brucellosis Seroprevalence on a Shared Winter Range. *The Journal of Wildlife Management* **66**:1236-1254.
- Gross, J. E., B. C. Lubow, and M. W. Miller. 2002. Modeling the epidemiology of brucellosis in the Greater Yellowstone Area. Pages 24-37 *in* T. J. Kreeger, editor. *Brucellosis in elk and bison in the Greater Yellowstone Area*. Wyoming Game and Fish Dept., Cheyenne, WY.
- Hamlin, K. L. 2006. Monitoring and Assessment of Wolf-Ungulate Interactions and Population Trends within the Greater Yellowstone Area, Southwestern Montana, and Montana Statewide. *Montana Fish, Wildlife & Parks*.
- Hamlin, K. L., and J. A. Cunningham. 2008. Montana elk movements, distribution, and numbers relative to brucellosis transmission risk. *Montana Fish, Wildlife, and Parks*.
- Hijmans, R. J., and J. Van Etten. 2010. raster: Geographic analysis and modeling with raster data. R package version 1.0.4. <http://CRAN.R-project.org/package=raster>.
- Hobbs, N. T., R. Wallen, J. Treanor, C. Geremia, and P. J. White. 2009. A stochastic population model of the Yellowstone bison population. National Park Service, Yellowstone Center for Resources, Yellowstone National Park, Wyoming.

- Johnson, D. E. 1951. Biology of the elk calf, *Cervus canadensis nelsoni*. Journal of Wildlife Management **15**:396-410.
- Kilpatrick, A. M., C. M. Gillin, and P. Daszak. 2009. Wildlife-livestock conflict: the risk of pathogen transmission from bison to cattle outside Yellowstone National Park. Journal of Applied Ecology **46**:476-485.
- Kreeger, T. J., W. E. Cook, W. H. Edwards, P. H. Elzer, and S. C. Olsen. 2002. *Brucella abortus* strain RB51 vaccination in elk II. Failure of high dosage to prevent abortion. Journal of Wildlife Diseases **38**:27-31.
- Maichak, E. J., B. M. Scurlock, J. D. Rogerson, L. L. Meadows, A. E. Barbknecht, W. H. Edwards, and P. C. Cross. 2009. Effects of management, behavior, and scavenging on risk of brucellosis transmission in elk of western Wyoming. Journal of Wildlife Diseases **45**:398-410.
- Meagher, M. 1973. The bison of Yellowstone National Park. National Park Service, Washington, D.C., USA.
- Meagher, M., and M. E. Meyer. 1994. On the origin of brucellosis in bison of Yellowstone National Park: A review. Conservation Biology **8**:645-653.
- Olsen, S., A. E. Jensen, W. C. Stoffregen, and M. V. Palmer. 2003. Efficacy of calfhood vaccination with *Brucella abortus* strain RB51 in protecting bison against brucellosis. Research in Veterinary Science **74**:17-22.
- Peterson, M. J., W. E. Grant, and D. S. Davis. 1991. Bison-Brucellosis Management - Simulation of Alternative Strategies. Journal of Wildlife Management **55**:205-213.
- Plumb, G., and K. Aune. 2002. The long term Interagency Bison Management Plan for Yellowstone National Park and the State of Montana. Pages 136-145 in T. J. Kreeger,

- editor. Brucellosis in elk and bison in the Greater Yellowstone Area. Wyoming Department of Game and Fish, Cheyenne, WY.
- Proffitt, K. M., P. J. White, and R. A. Garrott. 2010. Spatio-temporal overlap between Yellowstone bison and elk -- implications of wolf restoration and other factors for brucellosis transmission risk. *Journal of Applied Ecology* **47**:281-289.
- R Development Core Team. 2010. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Roffe, T. J., J. C. Rhyan, K. Aune, L. M. Philo, D. R. Ewalt, T. Gidlewski, and S. G. Hennager. 1999. Brucellosis in Yellowstone National Park bison: Quantitative serology and infection. *Journal of Wildlife Management* **63**:1132-1137.
- Schumaker, B. A., J. A. K. Mazet, J. Treanor, R. Wallen, A. W. Tam, I. A. Gardner, M. Zaluski, and T. E. Carpenter. 2010. Bison or elk: Who should be the target of brucellosis control in the northern greater Yellowstone area? Submitted to *Journal of Applied Ecology*.
- Thorne, E. T., J. K. Morton, F. M. Blunt, and H. A. Dawson. 1978. Brucellosis in elk. II. Clinical effects and means of transmission as determined through artificial infections. *Journal of Wildlife Diseases* **14**:280-291.
- Treanor, J., J. S. Johnson, R. L. Wallen, S. Cilles, P. H. Crowley, J. J. Cox, D. S. Maehr, P. J. White, and G. E. Plumb. 2010. Vaccination strategies for managing brucellosis in Yellowstone bison. *Vaccine* **28S**:F64-F72.
- USDI-NPS. 2010. Draft Environmental Impact Statement for Brucellosis Remote Vaccination Program for Bison in Yellowstone National Park. US Department of Interior.
- Yellowstone Center for Resources. 2008. Vaccination strategies for managing brucellosis in Yellowstone bison. National Park Service, Yellowstone National Park, Wyoming.

Yellowstone Center for Resources. 2009. Parturition in Yellowstone bison. National Park Service, Mammoth Hot Springs, Wyoming.

Table 1. Input parameters for a *Brucella abortus* transmission model used to assess the risk of an infectious event occurring in elk and bison populations in the northern greater Yellowstone area.

Description of variables	Statistical distribution (parameters) [Mean, SD]	Source
Shedding proportion	Beta (12,14) [0.46, 0.10]	(Roffe et al. 1999) ^a
Fetal persistence	BetaGeneral (2, 6.93, 1, 78) [18.25, 10.19]	(Aune et al. 2007)
<u>Bison</u>		
Number of animals Fit from 2000-2008 data	Logistic (3788.53, 450.13) [3788.53, 816.45]	(National Park Service, unpublished data)
Age proportion (of total population): Fit from 2004-2008 data		(National Park Service, unpublished data)
2-3 year-old females	BetaSubjective (0.043, 0.047, 0.04736, 0.053) [0.047, 0.002]	
4+ year-old females	Pareto (46.43, 0.35123) [0.36, 0.01]	
Proportion pregnant:		
2-3 year-old	Uniform (0.71, 0.79) [0.75, 0.02]	(Yellowstone Center for Resources 2008)
4+ year-old	Uniform (0.76, 0.89) [0.83, 0.04]	
Proportion seropositive 2+ year-old <u>(sampled at boundary capture facility)</u>	Beta (331.0, 211.6) [0.61, 0.02]	(National Park Service, unpublished data)

Percentage shedding by abortion:

First pregnancy females	BetaSubjective (0.65, 0.78, 0.78, 0.9) [0.78, 0.07]	(Davis et al. 1990)
Mature females	BetaSubjective (0.01, 0.1, 0.09, 0.15) [0.09, 0.03]	(Peterson et al. 1991)
Birth synchrony	Normal (40.57, 13.33) [40.57, 13.33] Day 1 = April 1	(Berger and Cain 1999)

Elk

Adult female proportion Fit from 2000-2008 data	BetaSubjective (0.52, 0.73, 0.7, 0.8) [0.7, 0.06]	(National Park Service, unpublished data)
Adult female: yearling	10:1	(National Park Service, unpublished data)
Proportion pregnant: Fit from 2000-2006 data		(National Park Service, unpublished data)
Yearling	BetaSubjective (0.1, 0.33, 0.32, 0.4) [0.32, 0.03]	
Adult	BetaSubjective (0.78, 0.82, 0.815, 0.84) [0.82, 0.01]	
Percentage of shedding by abortion:		
First pregnancy females	Beta (13.3, 14.4) [0.48, 0.09]	(Thorne et al. 1978)
Mature females	Beta (1.2, 6.8) [0.15, 0.12]	
Birth synchrony	Poisson (32.526) [32.526, 5.703] Day 1 = May 1	(Maichak et al. 2009)

Gallatin-Madison

Number of animals	Normal (7807, 793)	
Fit from 2000-2008 estimates (sightability corrected using 1.322 correction factor)	[7807, 793]	(Hamlin and Cunningham 2008)

B. abortus seropos. proportion Beta (3.1, 101.5)
[0.03, 0.02]

Gravelly-Snowcrest

(Hamlin 2006)

Number of animals Uniform (10,900, 11,570)
Fit from 2004&2006 data [11,235, 193]

B. abortus seropos. proportion Beta (3.1, 101.5)
[0.03, 0.02]

Madison-Firehole

Number of animals Loglogistic (236.8, 196.2, 1.4)
Fit from 2000-2008 estimates [757.2, N/A] (Hamlin and Cunningham 2008)
(sightability corrected using 1.322 correction factor)

B. abortus seropos. proportion Beta (3.1, 101.5)
[0.03, 0.02]

Northern Yellowstone

Number of animals Lognormal (9742, 3801, Shift (3396))
Fit from 2000-2008 estimates [13,137, 3800] (Cross et al. 2009)
(sightability corrected using 1.322 correction factor)

B. abortus seropos. proportion Uniform (0.01, 0.05) (Barber-Meyer et al. 2007)
[0.03, 0.01]

Sand Creek, Idaho

Number of adult females, 2006 1,413 (Mark Drew, Idaho
(sightability corrected using 1.322 correction factor) Department of Fish and
Game, unpublished data)

B. abortus seropos. proportion Beta (0.9, 100)
[0.01, 0.01]

a – study generalizes statistic for seropositive female bison

Table 2. Risk equation matrix for inter- and intra-species wildlife *Brucella abortus* transmission risk within and between bison and elk populations in the northern greater Yellowstone area.

***B. abortus* transmission risk equations**

Population at-risk	From Bison Population	From other Elk Population	From own Elk Population
Bison	(1)	(2)	N/A
<u>Elk Populations</u>			
Gallatin-Madison (GM)	(3)	(4)	(5)
Gravelly-Snowcrest (GS)	(6)	(7)	(8)
Madison-Firehole (MF)	(9)	(10)	(11)
Northern Yellowstone (NY)	(12)	(13)	(14)
Sand Creek, Idaho (SC)	(15)	(16)	(17)

(1) – (Bison shedding) * λ

(2) – (Elk shedding overlap) * γ

(3) – (Bison shedding overlap with GM) * δ

(4) – (GS, MF, NY, SC shedding overlap with GM) * ϵ

- (5) – (GM shedding) * θ
- (6) – (Bison shedding overlap with GS) * δ
- (7) – (GM, MF, NY, SC shedding overlap with GS) * ε
- (8) – (GS shedding) * θ
- (9) – (Bison shedding overlap with MF) * δ
- (10) – (GM, GS, NY, SC shedding overlap with MF) * ε
- (11) – (MF shedding) * θ
- (12) – (Bison shedding overlap with NY) * δ
- (13) – (GM, GS, MF, SC shedding overlap with NY) * ε
- (14) – (NY shedding) * θ
- (15) – (Bison shedding overlap with SC) * δ
- (16) – (GM, GS, MF, NY shedding overlap with SC) * ε
- (17) – (SC shedding) * θ

Probabilities of adequate contact, given spatiotemporal overlap

λ – Bison from bison

γ – Bison from elk

δ – Elk from bison


ε – Elk from other elk population

θ – Elk from own population

Table 3. Median risk, percentage of total, and 95% probability interval (P.I.) of *Brucella abortus* risk of transmission within and between bison and elk populations in the GYA using home range estimates for average winters. Units of risk are female exposure event-days.

Population at-risk	<i>B. abortus</i> transmission risk; % of total risk (95% P.I.)		
	From Bison Population	From other Elk Population	From own Elk Population
Bison	91.1; 78.3 (22.7, 242.4)	25.2; 21.7 (4.5, 98.5)	N/A
<u>Elk Populations</u>			
Gallatin-Madison	0.01; 0.2 (0.003, 0.03)	0.4; 7.7 (0.1, 1.3)	4.8; 92.1 (0.9, 18.0)
Gravelly-Snowcrest	0.002; <0.1 (0.001, 0.006)	0.06; 0.9 (0.01, 0.22)	6.9; 99.1 (1.3, 25.8)
Madison-Firehole	0.006; 0.5 (0.001, 0.016)	0.8; 72.3 (0.2, 3.1)	0.3; 27.1 (0.05, 1.8)
Northern Yellowstone	0.03; 0.4 (0.007, 0.09)	0.02; 0.3 (0.003, 0.06)	7.8; 99.4 (1.4, 31.5)
Sand Creek, Idaho	0.001; 0.2 (0, 0.001)	0.3; 59.9 (0.05, 1.0)	0.2; 39.9 (0.01, 1.6)

Figure 1. Map of bison and elk population distributions in the northern portion of the greater Yellowstone area based on an average winter.



Equation 1. Risk equation for inter- and intra-species wildlife *Brucella abortus* transmission risk model

Risk \propto

(Number of number of animals in the at-risk population) \times (Seronegative female proportion in the at-risk population) \times (Number of infectious event-days from the source of risk) \times (Proportion of shedding events overlapped by at-risk population) \times (Proportion of at-risk population exposed to risk source) \times Pr (Adequate Contact | Overlap)[†]

[†] - Probability of contact which would result in transmission if the exposed animal was susceptible, given overlap occurs

Equation 2. Incident Cases = (Prevalent Animals) / (Median Duration of Infection)^a

Objective 3

BRUCELLOSIS MANAGEMENT SCENARIOS IN THE NORTHERN GREATER YELLOWSTONE AREA

**Brant A. Schumaker,^{1*} Jonna A.K. Mazet,² John Treanor,³ Rick Wallen,³ Ian A. Gardner,⁴
Martin Zaluski,⁵ and Tim Carpenter¹**

¹Center for Animal Disease Modeling and Surveillance (CADMS), Department of Medicine and Epidemiology, School of Veterinary Medicine, University of California, Davis, CA 95616, USA;

²Wildlife Health Center, School of Veterinary Medicine, University of California, Davis, CA 95616, USA; ³National Park Service, Yellowstone National Park, P.O. Box 168, Wyoming

82190, USA; ⁴Department of Medicine and Epidemiology, School of Veterinary Medicine,

University of California, Davis, CA 95616, USA; ⁵Montana Department of Livestock, P.O. Box 202001, Helena, MT 59620-2001, USA;

*Corresponding author (email: theschu@ucdavis.edu, phone: (530) 752-3566, fax: (530) 752-1618)

Abstract

Bison (*Bison bison*) and elk (*Cervus elaphus*) in the northern portion of the greater Yellowstone area (GYA) remain a source for *Brucella abortus* infection in livestock. To increase tolerance for bison outside Yellowstone National Park (YNP) and reduce risk of cattle *B. abortus* exposure, the National Park Service has been exploring the option for the remote delivery of the RB51 brucellosis vaccine to various segments of the YNP bison herd. The parameters of a previously developed risk model were modified to evaluate the relative benefits of various management strategies to reduce wildlife *B. abortus* transmission to cattle on public grazing allotments in the northern GYA. Bison vaccination did not meaningfully reduce *B. abortus* transmission risk to cattle. Effective strategies included delaying the turn-on date to cattle grazing allotments, reducing elk seroprevalence, reducing the number of cattle at-risk, or prohibiting the comingling of elk and cattle on individual premises. The benefits of a later turn-on date for susceptible cattle provide perhaps the easiest method for reducing the incidence of livestock *B. abortus* infection in the GYA. Combining this strategy with best management practices to reduce comingling on individual premises will reduce the occurrence of cattle brucellosis in the northern GYA.

Keywords: bison, Brucella abortus, disease modeling, elk, emerging infectious disease, population management, RB51, risk management risk modeling, wildlife disease

INTRODUCTION

B. abortus is a gram-negative, facultative, intracellular bacterium that causes disease in many domestic and wild animal species including cattle, bison (*Bison bison*), and elk (*Cervus elaphus*) (Creech 1930, Thorne et al. 1978). Bacteria invade the mucous membranes of ungulates and can cause placentitis with late-gestation abortions in females and orchitis and epididymitis in males (Bercovich 1998). Increased abortion rates, decreased milk production, loss of condition, infertility, and lameness in cattle have made brucellosis extremely important to beef and milk producers around the world (Manthei and Carter 1950), restricting international trade in many instances (Wilson and Beers 2001). The bacterium can also be transmitted to humans as perhaps the most common zoonotic disease worldwide (Pappas et al. 2006)

The eradication of the disease from the United States has been a priority of the federal government since 1934, when a cooperative state-federal brucellosis eradication program (BEP) was adopted to reduce the prevalence of brucellosis in cattle, designating it the most significant livestock disease at that time. Since then, agencies have implemented a variety of livestock, wildlife, and disease risk management strategies (Cheville et al. 1998). Billions of dollars have been spent eradicating brucellosis from livestock in nearly every state in the US (Wise 1980). During the 76-year history of the BEP, it has limited the impact of brucellosis in cattle throughout the United States (Donch and Gertonson 2008). By early 2008, the United States and associated territories were all brucellosis free in livestock. However, in June 2008 brucellosis was again detected in cattle herds in Montana and Wyoming. Incidents in the last four years in all three states surrounding Yellowstone National Park (YNP) – Idaho, Montana, and Wyoming – have highlighted the importance of wildlife brucellosis.

Brucellosis was first detected among wildlife in and around YNP in 1917, when epizootic abortion was described in Yellowstone bison (Mohler, 1917). The disease was most likely acquired from domestic cattle, which were brought into the area for grazing (Meagher and Meyer 1994). Today, elk populations in the northern GYA have low seroprevalence (i.e., exposure; <5%) for *B. abortus*, whereas seroprevalence in Yellowstone bison is high (40-60%) (Cheville et al. 1998). Bison conservation continues to be a priority of the National Park Service; however, for decades, livestock and regulatory personnel have viewed Yellowstone bison as the primary source of *B. abortus* transmission risk to cattle because of their higher seroprevalence (Meagher and Meyer 1994). However, current management, which maintains spatial and temporal separation between bison and cattle, makes the risk of *B. abortus* transmission from bison to cattle in the northern GYA negligible (Kilpatrick et al., 2009(Schumaker et al. 2010)). However, hazing and culling actions by bison managers to maintain this separation have been highly scrutinized and criticized for their economic costs and negative effects to bison. In the last decade, there have been multiple detections of brucellosis in cattle in the GYA states (Idaho, Montana, Wyoming), with elk identified as the source of infection for nine cases since 2002 (Donch and Gertonson 2008).

The Interagency Brucellosis Management Plan (IBMP) was established in 2000 to manage the risk of *B. abortus* transmission from bison to cattle by implementing hazing, test-and-slaughter, hunting, and other actions near the boundary of Yellowstone National Park (Plumb and Aune 2002, Donch et al. 2005). These actions have successfully prevented the transmission of *B. abortus* from bison to cattle (Clarke et al. 2005), and an assessment suggest the risk of future *B. abortus* transmission is minimal under current management conditions (Kilpatrick et al. 2009). Since 2000, about 3,200 bison have been removed from the Yellowstone

herd with over 1000 animals, or 20% of the total population culled during the winter of 2005-2006. These actions have been controversial with animal advocacy groups.

Since the early period of the BEP, vaccination has been considered as a control method for *B. abortus* transmission. Because the serologic cross-reactions of strain 19 make it ineffective for test-and-cull methods of *B. abortus* control, other candidate vaccines were explored. A live rifampin-resistant “rough”, or devoid of the LPS O-chain, attenuated strain of *B. abortus* labeled “51” by internal laboratory nomenclature was developed by Schurig and colleagues (1991) and was later trademarked by Virginia Tech Intellectual Properties in 1992. Rough *Brucella* 51 (RB51) has proven to be less abortigenic in cattle than S19 while showing similar efficacy. Because RB51 lacks the O-chain on its LPS it does not cross-react on *B. abortus* serologic tests. Age-specific seroprevalence proportions in Yellowstone bison indicate that approximately 50% of bison are exposed prior to reproductive maturity (Treanor et al. 2007). Thus, early exposure to the vaccine may allow immature bison to develop resistance to infection, which could be maintained by booster vaccinations to reduce the occurrence of *B. abortus*-induced abortions.

To increase tolerance for bison outside YNP and reduce risk of cattle *B. abortus* exposure, the National Park Service has been exploring the option for the remote delivery of the RB 51 brucellosis vaccine to various segments of the YNP bison herd (USDI-NPS 2010). Vaccination of all female bison within YNP is expected to significantly reduce the population seroprevalence of *B. abortus* infection (Yellowstone Center for Resources 2008). However, a risk assessment of *B. abortus* transmission among elk, bison, and cattle in the northern portion of the GYA estimated the risk of bacterial transmission from bison to cattle under current management conditions to be minimal (Objective 1). Also, the likelihood of *B. abortus* transmission from bison to elk was shown to be minimal (Objective 2). The purpose of this study

was to evaluate meaningful *B. abortus* risk management alternatives to determine their relative efficacy in reducing *B. abortus* transmission risk from wildlife to cattle.

MATERIALS AND METHODS

Greater Yellowstone Area and Wildlife Populations

Yellowstone National Park (YNP) was established as America's first national park in 1872, and has become a flagship for wildlife conservation worldwide. Despite its large size of 8,987 square kilometers, YNP is not independent of its surrounding ecosystem, the greater Yellowstone area (GYA). The GYA is one of the largest intact temperate zone ecosystems on earth and includes approximately 28,000 square miles in Montana, Idaho and Wyoming and encompasses state lands, two national parks, portions of six national forests, three national wildlife refuges, Bureau of Land Management holdings, and private and tribal lands. The GYA is also home to the largest wild and free-ranging elk and bison populations in the United States.

The continental divide runs from west to east across the southern portion of YNP. The northern GYA includes the Yellowstone bison population and five elk populations (Gallatin-Madison, Gravelly-Snowcrest, Madison-Firehole, northern Yellowstone, and Sand Creek, Idaho), which are distributed across over 1,100 square miles in the northern GYA. Estimates of northern Yellowstone elk were near 25,000 animals in the late 1980's, but decreased by approximately 50-60% by 2006 (Eberhardt et al. 2007). The Yellowstone bison population ranges between 2000 and 5000 individuals (Meagher 1973, Clarke et al. 2005) depending on the season. The 2009 summer count for the Yellowstone bison herd was 3,300 animals, divided equally between a central and northern breeding population. These bison are desirable for the conservation of the species because the population is derived from the original wild herd and an

introduced herd containing widely diverse genetics (Meagher 1973). In addition, the bison have had no evidence of cattle-hybridization (Halbert et al. 2005). Therefore, disease management activities, including the future potential for movement of individual bison into other herds, are of special interest in this population.

Domestic cattle (266 in the winter and 1363 in the spring in 2006) are grazed on public and private lands adjacent to Yellowstone National Park (YNP) and within habitat occupied by bison and elk during the winter (Kilpatrick et al. 2009). Federal and state management agencies have attempted to decrease the risk of *B. abortus* transmission from bison to cattle using hazing and bison culling to maintain spatio-temporal separation from cattle (U.S. Department of Interior [USDI] and U.S. Department of Agriculture [USDA] 2000).

Risk model

The evaluation of efforts to reduce *B. abortus* transmission from wildlife to cattle in the northern GYA employed a previously developed risk model (Objective 1). The model estimated the number and spatiotemporal distribution of *B. abortus* shedding events from third-trimester abortions and infectious live parturition events from one bison and five elk populations in the northern GYA (Figure 1). The stochastic model was parameterized with statistical distributions fit to winter severity, animal location, serologic testing, demographic and epidemiologic data using @RISK v5.5 (Palisade Corporation, Ithaca, New York, USA; Table 1). The assumptions for the model were: 1) adult females are the primary source of infection; 2) the critical season of transmission is between January 1 and June 30; 3) no fully immune state exists; and 4) random mixing of animals occurs within a population.

Risk calculation

Risk of *B. abortus* transmission from wildlife to cattle is a combination of spatiotemporal overlap of at-risk individuals, the number and location of infectious events from the wildlife reservoirs. Numbers of infectious events in each wildlife population were taken from the results of 50,000 iterations of the stochastic risk model and distributed to fixed kernel density estimations of wildlife home ranges as described earlier (Objective 1). Overlap among elk and bison populations was calculated using the Spatial Analyst extension in ArcGIS v9.3 (Environmental Systems Research Institute, Redlands, California). Rasters were converted to ASCII files and the percentage of volume overlap was calculated using R statistical language v2.11.1 (R Development Core Team 2010) and the raster R package (Hijmans and Van Etten 2010).

Risk Management Alternatives

Risk management alternatives to reduce wildlife to cattle transmission of *B. abortus* in the northern portion of the GYA were explored. Because the study focused on the northern GYA, it did not explore the possibility of limiting or eliminating elk feedgrounds, which is a southern GYA issue. Also, strategies for early detection of cattle cases or to reduce disease spread within cattle populations were not examined, since the risk calculations were based on bacterial transmission from wildlife to cattle, not between cattle.

Strategy 1: Actively manage bison population between 2500-4500

The modeled food-limiting carrying capacity for bison within YNP is 6200 individuals (Plumb et al. 2009). However, even at lower population numbers, interactive effects of severe

winters and herd density with population numbers greater than 4200 have been associated with, and may contribute to, large-scale dispersal to lower elevations. Plumb et al. (2009), recommended the Yellowstone bison herd be maintained with less than 4500 animals to abate most large-scale movements outside the park during near-average winter conditions. Change to risk model parameters: constrain bison population distribution between 2500 and 4500.

Strategy 2: Reduction in wildlife population numbers

Multiple strategies have been suggested for reducing portions of the elk and bison populations in the northern GYA. The suggested strategies have included immunocontraception, hunting, or culling of seropositive animals. Change to risk model parameters: reduce individual wildlife population by a range of values (1, 5, 10, 20, 30% reduction).

Strategy 3: Bison vaccination

Treanor et al. (2010) concluded that vaccinating all Yellowstone female bison with RB51 would result in a reduction in seroprevalence from 47% to 16% over 30 years. Although the risk model started with a higher median seroprevalence, this strategy was modeled using the 30% absolute change in the seropositive proportion for bison. Change to risk model parameters: reduce median bison seropositive proportion from 0.61 to 0.30 and changed distribution to Beta (178.2, 415.8).

Strategy 4: Reduce elk seroprevalence

Elk have been vaccinated with strain 19 on the Wyoming feedgrounds. Also, a five-year pilot test-and-slaughter program around Pinedale, Wyoming by Laura Linn-Meadows lowered *B.*

abortus seroprevalence (USAHA, 2009). Removals of pre-reproductive seropositive elk are expected to reduce seroprevalence over time. Change to risk model parameters: reduce elk seroprevalence by a range of values (10, 20, 30, 50, 70% reduction).

Strategy 5: Reduce number of cattle at-risk

Reducing the number of susceptible cattle that graze on allotments within wildlife home ranges would decrease the risk of *B. abortus* transmission between wildlife and cattle. This could be accomplished by moving operations to non-overlapped allotments or switching the type of operation to avoid sexually-intact females in the herd. This could also be accomplished by vaccination of cattle. Change to risk model parameters: reduce cattle population by a range of values (10, 20, 30, 50, 70% reduction).

Strategy 6: Reduced tolerance for elk comingling with cattle

All recent detections of brucellosis in northern GYA cattle have been qualitatively attributed to elk that may or may not have seasonally occupied YNP (Galey et al., 2005). Due to the intense focus on bison *B. abortus* management during the past decade, elk have received minimal brucellosis management attention until recently and often move freely across the ecosystem and come into close contact with cattle premises. Best management practices include but are not limited to fencing the cattle feeding area and feed storage, altering feeding times, and hazing wildlife that are within close proximity to the feeding area or stack yard. Change to risk model parameters: incorporate the wildlife tolerance parameter previously used only for bison to the elk to cattle risk equation reducing risk. This was modeled as a Uniform (0,1) distribution.

Strategy 7: Delay earliest cattle turn-on date

Grazing on public lands in the northern greater Yellowstone area begins on different dates depending on the grazing allotment with an earliest “on date” of June 1st. Pushing back the earliest start date would reduce the number of days that each cow would be at-risk. Change to risk model parameters: set maximum days at-risk based on different earliest turn-on dates (6/2, 6/16, 6/21, 6/26, 6/30).

The parameters of the wildlife shedding model were altered to simulate the effects of each management strategy in order to evaluate their effects on *B. abortus* transmission risk. The risk model with new parameters was run for 50,000 iterations for each strategy evaluated and the median and 95% probability intervals (95% PIs) of total cattle exposure risk were determined. The risk was equated to the incidence of cattle brucellosis in the northern GYA during the study period.

RESULTS

The modeled baseline total risk to cattle in a year with a typical winter was 5.6 cattle-exposure infectious-event days (95% P.I. 0.2-55.3). Since there were two cases of *B. abortus* in cattle from 2004-2008, this risk calculation equated to 0.4 cases/year for cattle on grazing allotments. Therefore, it is estimated there were a median of 0.07 (0.4/5.6) infections/year for every annual cattle-exposure infectious event-day (95% P.I. 0.007-2.0), i.e. 1 in approximately every 14 cattle-exposure infectious-event days resulted in *B. abortus* transmission to cattle. The

results of the modeled management strategies showed variable success in lowering *B. abortus* transmission risk (Table 2). The goal for *B. abortus* risk management in the GYA is to reduce cattle infection incidence below the level that would result in a loss of *Brucella*-free status. The most frequent case incidence that would allow Montana to maintain its status is one case every three years. Active management to wildlife population numbers in the absence of reductions in *B. abortus* prevalence did not meaningfully reduce risk to cattle. A 30% reduction to the Gallatin-Madison elk population showed the largest effect on risk, however not enough for Montana to consistently stay *Brucella*-free. Bison vaccination and the resultant lowering of bison seroprevalence also was not an effective risk reduction strategy. Equal proportional reductions to elk *B. abortus* prevalence and the number of cattle at-risk showed equal benefit in reducing risk. Reductions in these parameters of 20% and 70% lowered the incidence to one case in 3.2 and 8.2 years, respectively. Reducing tolerance to elk comingling with cattle effectively reduced *B. abortus* incidence to one case every 6.8 years. Delaying cattle grazing turn-on dates showed the largest absolute reduction in *B. abortus* transmission incidence between wildlife and cattle. Delay the earliest cattle turn-on date to 6/21 reduced the transmission incidence to one case every 3.3 years and further delaying the turn-on date to 6/30 resulted in only one case every 30.4 years.

DISCUSSION

Disease management at the wildlife-livestock interface is hampered by the challenge of balancing wildlife conservation with the livelihoods and traditions of livestock producers. The potential for disease transmission between wildlife and livestock exacerbates conflicts between natural resource managers and cattlemen, reduces tolerance for wildlife near livestock

operations, and negatively impacts conservation. Therefore, diseases that affect both wildlife and livestock are important in resource management, regardless of their direct impact to the wild animal populations, which may serve as their reservoirs.

While the grazing season in the GYA is short, the benefits of a later turn-on date for susceptible cattle provide perhaps the easiest method for reducing the incidence of livestock *B. abortus* infection in the GYA. Combining this strategy with best management practices to reduce comingling on individual premises will make meaningful progress toward the eventual eradication of *B. abortus* from the US. Although, vaccination is expected to substantially reduce the prevalence of *B. abortus* in bison, with currently mandated separation of bison and cattle, it is unlikely to meaningfully reduce direct or indirect *B. abortus* transmission risk, from bison or elk, respectively. However, management alternatives that reduce *B. abortus* prevalence in bison would likely increase the tolerance for bison outside YNP boundaries, however, as they would decrease either the actual potential for transmission from bison to cattle or the perceived potential for transmission. Moving forward efforts to further reduce risk cattle must focus on either reducing *B. abortus* prevalence in elk or taking steps to locally reduce comingling of elk with susceptible cattle. The results of our analysis can be used to quantify the relative benefits of alternative management strategies in order for future policy to be based on objective science.

ACKNOWLEDGEMENTS

Funding for this project was provided by the US Department of Agriculture: Animal and Plant Health Inspection Service (APHIS) and the US National Park Service through the Yellowstone Wildlife Health Program. The project was also funded through the Foreign Animal and Zoonotic Disease Defense Center of Excellence, by a grant from the Department of Homeland Security,

Science and Technology Directorate, Office of University Programs. We also thank Glenn Plumb and P.J. White for pursuing funding and building the collaborative relationships that made this work possible.

REFERENCES

- Aune, K., J. C. Rhyan, B. Corso, and T. Roffe. 2007. Environmental persistence of *Brucella* organisms in natural environments of the greater Yellowstone area - A preliminary analysis. Report of the Committee on Brucellosis. United States Animal Health Association, Richmond, Virginia.
- Barber-Meyer, S. M., P. J. White, and L. D. Mech. 2007. Survey of selected pathogens and blood parameters of northern Yellowstone elk: wolf sanitation effect implications. *American Midland Naturalist* **158**:369-381.
- Bercovich, Z. 1998. Maintenance of *Brucella abortus*-free herds: A review with emphasis on the epidemiology and the problems in diagnosing brucellosis in areas of low prevalence. *Veterinary Quarterly* **20**:81-88.
- Berger, J., and S. L. Cain. 1999. Reproductive synchrony in brucellosis-exposed bison in the southern greater Yellowstone ecosystem and in noninfected populations. *Conservation Biology* **13**:357-366.
- Cheville, N., D. R. McCullough, and L. R. Paulson. 1998. Brucellosis in the greater Yellowstone area. National Research Council, Washington D.C.
- Clarke, R., C. Jourdonnais, J. Munding, L. Stoeffler, and R. Wallen. 2005. A Status Review of Adaptive Management Elements 2000 to 2005. Interagency Bison Management Plan.
- Creech, G. T. 1930. *Brucella abortus* infection in a male bison. *North American Veterinarian* **11**:35-36.

- Cross, P. C., T. O. Lemke, P. J. White, and D. B. Tyers. 2009. Northern Yellowstone cooperative wildlife working group 2008 annual report (October 1, 2007-September 30, 2008). U.S. Geological Survey, Northern Rocky Mountain Science Center, Bozeman, Montana.
- Davis, D., J. W. Templeton, T. A. Ficht, J. D. Williams, J. D. Kopec, and L. G. Adams. 1990. *Brucella abortus* in captive bison. I. Serology, bacteriology, pathogenesis, and transmission to cattle. *Journal of Wildlife Diseases* **26**:360-371.
- Davis, D. S., and P. H. Elzer. 2002. *Brucella* vaccines in wildlife. *Veterinary Microbiology* **90**:533-544.
- Donch, D. A., and A. A. Gertonson. 2008. Status report -- Fiscal year 2008; Cooperative State-Federal Brucellosis Eradication Program. USDA-APHIS Veterinary Services.
- Donch, D. A., A. A. Gertonson, J. C. Rhyhan, and M. J. Gilsdorf. 2005. Status Report - Fiscal Year 2005; Cooperative State-Federal Brucellosis Eradication Program. USDA-APHIS Veterinary Services.
- Eberhardt, L. L., P. J. White, R. A. Garrott, and D. B. Houston. 2007. A seventy-year history of trends in Yellowstone's northern elk herd. *Journal of Wildlife Management* **71**:594-602.
- Ferrari, M. J., and R. A. Garrot. 2002. Bison and Elk: Brucellosis Seroprevalence on a Shared Winter Range. *The Journal of Wildlife Management* **66**:1236-1254.
- Halbert, N. D., T. J. Ward, R. D. Schnabel, J. F. Taylor, and J. N. Derr. 2005. Conservation genomics: Disequilibrium mapping of domestic cattle chromosomal segments in North American bison populations. *Molecular Ecology* **14**:2343-2362.
- Hamlin, K. L. 2006. Monitoring and Assessment of Wolf-Ungulate Interactions and Population Trends within the Greater Yellowstone Area, Southwestern Montana, and Montana Statewide. Montana Fish, Wildlife & Parks.

- Hamlin, K. L., and J. A. Cunningham. 2008. Montana elk movements, distribution, and numbers relative to brucellosis transmission risk. *Montana Fish, Wildlife, and Parks*.
- Hijmans, R. J., and J. Van Etten. 2010. raster: Geographic analysis and modeling with raster data. R package version 1.0.4. <http://CRAN.R-project.org/package=raster>.
- Johnson, D. E. 1951. Biology of the elk calf, *Cervus canadensis nelsoni*. *Journal of Wildlife Management* **15**:396-410.
- Jones, J. D., J. J. Treanor, and R. L. Wallen. 2009. Parturition in Yellowstone bison. National Park Service, Yellowstone National Park, Wyoming.
- Kilpatrick, A. M., C. M. Gillin, and P. Daszak. 2009. Wildlife-livestock conflict: the risk of pathogen transmission from bison to cattle outside Yellowstone National Park. *Journal of Applied Ecology* **46**:476-485.
- Kreeger, T. J., W. E. Cook, W. H. Edwards, P. H. Elzer, and S. C. Olsen. 2002. *Brucella abortus* strain RB51 vaccination in elk II. Failure of high dosage to prevent abortion. *Journal of Wildlife Diseases* **38**:27-31.
- Maichak, E. J., B. M. Scurlock, J. D. Rogerson, L. L. Meadows, A. E. Barbknecht, W. H. Edwards, and P. C. Cross. 2009. Effects of management, behavior, and scavenging on risk of brucellosis transmission in elk of western Wyoming. *Journal of Wildlife Diseases* **45**:398-410.
- Manthei, C. A., and R. W. Carter. 1950. Persistence of *Brucella abortus* infection in cattle. *American Journal of Veterinary Research* **11**:173-180.
- Meagher, M. 1973. The bison of Yellowstone National Park. National Park Service, Washington, D.C., USA.

- Meagher, M., and M. E. Meyer. 1994. On the origin of brucellosis in bison of Yellowstone National Park: A review. *Conservation Biology* **8**:645-653.
- Olsen, S., A. E. Jensen, W. C. Stoffregen, and M. V. Palmer. 2003. Efficacy of calfhood vaccination with *Brucella abortus* strain RB51 in protecting bison against brucellosis. *Research in Veterinary Science* **74**:17-22.
- Olsen, S. C., R. J. Christie, D. W. Grainger, and W. S. Stoffregen. 2006. Immunologic responses of bison to vaccination with *Brucella abortus* strain RB51: comparison of parenteral to ballistic delivery via compressed pellets or photopolymerized hydrogels. *Vaccine* **24**:1346-1353.
- Pappas, G., P. Papadimitriou, N. Akritidis, L. Christou, and E. Tsianos. 2006. The new global map of human brucellosis. *Lancet Infectious Diseases* **6**:91-99.
- Peterson, M. J., W. E. Grant, and D. S. Davis. 1991. Bison-Brucellosis Management - Simulation of Alternative Strategies. *Journal of Wildlife Management* **55**:205-213.
- Plumb, G., and K. Aune. 2002. The long term Interagency Bison Management Plan for Yellowstone National Park and the State of Montana. Pages 136-145 in T. J. Kreeger, editor. *Brucellosis in elk and bison in the Greater Yellowstone Area*. Wyoming Department of Game and Fish, Cheyenne, WY.
- Plumb, G. E., P. J. White, M. B. Coughenour, and R. L. Wallen. 2009. Carrying capacity, migration, and dispersal in Yellowstone bison. *Biological Conservation* **142**:2377-2387.
- Proffitt, K. M., P. J. White, and R. A. Garrott. 2010. Spatio-temporal overlap between Yellowstone bison and elk -- implications of wolf restoration and other factors for brucellosis transmission risk. *Journal of Applied Ecology* **47**:281-289.

- R Development Core Team. 2010. R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria.
- Roffe, T. J., L. C. Jones, K. Coffin, M. L. Drew, S. J. Sweeney, S. D. Hagius, P. H. Elzer, and D. Davis. 2004. Efficacy of single calfhooed vaccination of elk with *Brucella abortus* strain 19. *Journal of Wildlife Management* **68**:830-836.
- Roffe, T. J., J. C. Rhyan, K. Aune, L. M. Philo, D. R. Ewalt, T. Gidlewski, and S. G. Hennager. 1999. Brucellosis in Yellowstone National Park bison: Quantitative serology and infection. *Journal of Wildlife Management* **63**:1132-1137.
- Schumaker, B. A., J. A. K. Mazet, J. Treanor, R. Wallen, A. W. Tam, I. A. Gardner, M. Zaluski, and T. E. Carpenter. 2010. Bison or elk: Who should be the target of brucellosis control in the northern greater Yellowstone area? Submitted to *Journal of Applied Ecology*.
- Schurig, G. G., R. M. Roop II, T. Bagchi, S. Boyle, D. Buhrman, and N. Sriranganathan. 1991. Biological properties of RB51; a stable rough strain of *Brucella abortus*. *Veterinary Microbiology* **28**:171-188.
- Thorne, E. T., J. K. Morton, F. M. Blunt, and H. A. Dawson. 1978. Brucellosis in elk. II. Clinical effects and means of transmission as determined through artificial infections. *Journal of Wildlife Diseases* **14**:280-291.
- Treanor, J., J. Johnson, R. Wallen, S. Cilles, P. Crowley, and D. Maehr. 2008. Vaccination strategies for managing brucellosis in Yellowstone bison. National Park Service, Yellowstone National Park, Wyoming.
- Treanor, J. J., J. S. Johnson, R. L. Wallen, S. Cilles, P. H. Crowley, J. J. Cox, D. S. Maehr, P. J. White, and G. E. Plumb. 2010. Vaccination strategies for managing brucellosis in Yellowstone bison. *Vaccine* **28S**:F64-F72.

- U.S. Department of Interior [USDI], and U.S. Department of Agriculture [USDA]. 2000. Record of Decision for Final Environmental Impact Statement and Bison Management Plan for the State of Montana and Yellowstone National Park. USDI National Park Service and USDA Animal and Plant Health Inspection Service.
- USDI-NPS. 2010. Draft Environmental Impact Statement for Brucellosis Remote Vaccination Program for Bison in Yellowstone National Park. US Department of Interior.
- Wilson, D. W., and P. T. Beers. 2001. Global trade requirements and compliance with World Trade Organization agreements: the role of tracing animals and animal products. *Revue Scientifique et Technique Office International des Epizooties* **20**:379-384.
- Wise, R. I. 1980. Brucellosis in the United States: Past, present, and future. *Journal of the American Medical Association* **244**:2318-2322.
- Yellowstone Center for Resources. 2008. Vaccination strategies for managing brucellosis in Yellowstone bison. National Park Service, Yellowstone National Park, Wyoming.
- Yellowstone Center for Resources. 2009. Parturition in Yellowstone bison. National Park Service, Mammoth Hot Springs, Wyoming.

Table 1. Baseline input parameters for a *Brucella abortus* transmission model used to assess the risk of an infectious event occurring in elk and bison populations in the northern greater Yellowstone area.

Description of variables	Statistical distribution (parameters) [Mean, SD]	Source
Shedding proportion	Beta (12,14) [0.46, 0.10]	(Roffe et al. 1999) ^a
Fetal persistence	BetaGeneral (2, 6.93, 1, 78) [18.25, 10.19]	(Aune et al. 2007)
<u>Bison</u>		
Number of animals Fit from 2000-2008 data	Logistic (3788.53, 450.13) [3788.53, 816.45]	(National Park Service, unpublished data)
Age proportion (of total population): Fit from 2004-2008 data		(National Park Service, unpublished data)
2-3 year-old females	BetaSubjective (0.043, 0.047, 0.04736, 0.053) [0.047, 0.002]	
4+ year-old females	Pareto (46.43, 0.35123) [0.36, 0.01]	
Proportion pregnant: 2-3 year-old	Uniform (0.71, 0.79) [0.75, 0.02]	(Yellowstone Center for Resources 2008)
4+ year-old	Uniform (0.76, 0.89) [0.83, 0.04]	
Proportion seropositive 2+ year-old (sampled at boundary capture facility)	Beta (331.0, 211.6) [0.61, 0.02]	(National Park Service, unpublished data)
Percentage shedding by abortion: First pregnancy females	BetaSubjective (0.65, 0.78, 0.78, 0.9) [0.78, 0.07]	(Davis et al. 1990)

Mature females	BetaSubjective (0.01, 0.1, 0.09, 0.15)(Peterson et al. 1991) [0.09, 0.03]	
Birth synchrony	Normal (40.57, 13.33) [40.57, 13.33] Day 1 = April 1	(Berger and Cain 1999)

Elk

Adult female proportion Fit from 2000-2008 data	BetaSubjective (0.52, 0.73, 0.7, 0.8) [0.7, 0.06]	(National Park Service, unpublished data)
Adult female: yearling	10:1	(National Park Service, unpublished data)
Proportion pregnant: Fit from 2000-2006 data		(National Park Service, unpublished data)
Yearling	BetaSubjective (0.1, 0.33, 0.32, 0.4) [0.32, 0.03]	
Adult	BetaSubjective (0.78, 0.82, 0.815, 0.84) [0.82, 0.01]	
Percentage of shedding by abortion: First pregnancy females	Beta (13.3, 14.4) [0.48, 0.09]	(Thorne et al. 1978)
Mature females	Beta (1.2, 6.8) [0.15, 0.12]	
Birth synchrony	Poisson (32.526) [32.526, 5.703] Day 1 = May 1	(Maichak et al. 2009)

Gallatin-Madison

Number of animals Fit from 2000-2008 estimates [7807, 793] (sightability corrected using 1.322 correction factor)	Normal (7807, 793)	(Hamlin and Cunningham 2008)
<i>B. abortus</i> seropos. proportion	Beta (3.1, 101.5) [0.03, 0.02]	

Gravelly-Snowcrest

Number of animals	Uniform (10,900, 11,570)	(Hamlin 2006)
-------------------	--------------------------	---------------

Fit from 2004&2006 data [11,235, 193]
(sightability corrected using 1.322 correction factor)

B. abortus seropos. proportion Beta (3.1, 101.5)
[0.03, 0.02]

Madison-Firehole

Number of animals Loglogistic (236.8, 196.2, 1.4) (Hamlin and Cunningham 2008)
Fit from 2000-2008 estimates [757.2, N/A]
(sightability corrected using 1.322 correction factor)

B. abortus seropos. proportion Beta (3.1, 101.5)
[0.03, 0.02]

Northern Yellowstone

Number of animals Lognormal (9742, 3801, Shift (3396)) (Cross et al. 2009)
Fit from 2000-2008 estimates [13,137, 3800]
(sightability corrected using 1.322 correction factor)

B. abortus seropos. proportion Uniform (0.01, 0.05) (Barber-Meyer et al. 2007)
[0.03, 0.01]

Sand Creek, Idaho

Number of adult females, 2006 1,413 (Mark Drew, Idaho
(sightability corrected using 1.322 correction factor) Department of Fish and
Game, unpublished data)

B. abortus seropos. proportion Beta (0.9, 100)
[0.01, 0.01]


a – study generalizes statistic for seropositive female bison

Table 2. Median cattle risk of exposure to a *Brucella abortus* wildlife infectious shedding event and cattle infection incidence for a typical winter. Bolded strategies and results indicate a reduction in risk that would prevent the loss of *Brucella*-free status.

Strategy	Risk estimate (cattle-exposure infectious event-days)	Infection incidence ⁻¹ (years/case)
1. Actively maintain bison population between 2500-4500	5.6	2.5
2. Reduce wildlife population numbers (1, 5, 10, 20, 30% reduction)		
Bison	5.6, 5.6, 5.5, 5.5, 5.5	2.5, 2.5, 2.6, 2.6, 2.6
Gallatin-Madison elk	5.5, 5.4, 5.3, 5.1, 4.8	2.6, 2.6, 2.6, 2.8, 2.9
Gravelly-Snowcrest elk	5.5, 5.5, 5.4, 5.3, 5.1	2.6, 2.6, 2.6, 2.6, 2.8
Northern Yellowstone elk	5.6, 5.4, 5.4, 5.2, 5.0	2.5, 2.6, 2.6, 2.7, 2.8
3. Bison vaccination (Reduce bison seroprevalence from 61% to 30%).	5.5	2.6
4. Reduce elk seroprevalence (10, 20, 30, 50, 70% reduction)	5.0, 4.4, 3.9, 2.8, 1.7	2.8, 3.2, 3.6, 5.0, 8.2
5. Reduce number of cattle at-risk (10, 20, 30, 50, 70% reduction)	5.0, 4.4, 3.9, 2.8, 1.7	2.8, 3.2, 3.6, 5.0, 8.2
6. Reduce tolerance for elk comingling with cattle	2.1	6.8

7. Delay earliest cattle turn-on date (6/2, 6/16, 6/21 , 6/26 , 6/30)	5.5, 5.2, 4.3, 2.3, 0.5	2.6, 2.7, 3.3 , 6.1 , 30.4
---	-------------------------	---

Figure 1. Map of bison and elk population distributions in the northern portion of the greater Yellowstone area based on an average winter.



CONCLUSIONS

This research was initiated concurrently with the establishment of the Yellowstone Wildlife Health Program (YWHP). The YWHP, a cooperative partnership between Montana State University, the University of California, Davis, and YNP, was created to help answer meaningful scientific research questions and establish professional networks to funnel the answers to these questions back to YNP. An organizational workshop listed brucellosis among the highest priority research needs of YNP and identified risk assessments, transmission dynamics, and diagnostic test evaluations as specific scientific needs (Schumaker et al., 2007).

The study presented the first spatially-explicit framework for assessing the risk of bacterial shedding of *B. abortus* by bison and elk across the northern portion of the GYA. Although our results support substantial shedding of *B. abortus* from bison in some winters, the most substantial risk of *B. abortus* transmission to cattle was from elk. Future risk estimates for bison depend on adaptive management of the population. Interactive effects between population size and winter severity were major determinants influencing bison movements to lower elevation winter grazing areas and overlap with federally-regulated domestic cattle grazing allotments. However, during the critical period of potential *B. abortus* exposure to cattle, the risk from Yellowstone bison was minimal. Natural movements of animals back to higher elevation summer ranges and boundary management operations were important in minimizing the contribution of bison to cattle exposure risk, which supports continued boundary management operations for spatio-temporal separation between bison and cattle. Under current management practices, bison risk to cattle grazing in the northern portion of the GYA is expected to remain small.

In addition to spatio-temporal overlap of wildlife home ranges and cattle grazing allotments, the major contributors to risk were wildlife population size and the number of elk that were shedding *B. abortus*. While elk currently have a lower density of shedding events throughout their range, they have a larger overlap with cattle and are more tolerated by managers and livestock keepers on public grazing allotments. With increased disease prevalence due to increased winter densities or other factors, elk will likely contribute greatly to the overall level of bacterial shedding on the northern GYA landscape and represent the vast majority of risk of *B. abortus* exposure to cattle grazing in the northern portion of the GYA. Therefore, brucellosis management efforts should increasingly focus on the comingling of cattle and elk during the critical abortion period to more effectively decrease risk of transmission.

Continued exploration of the brucellosis risk model found that minimal opportunity exists for *B. abortus* transmission from bison to elk under natural conditions in the northern GYA. The reasons for this lower probability of adequate contact for *B. abortus* transmission, even when spatio-temporal overlap occurred, are likely immunological or behavioral. The risk model may be expanded to include the entire GYA or serve as a template for models of other diseases. As additional data become available, especially additional spatial locations of cattle and wildlife and animal movement information, the model can be refined for even more targeted management decisions. Current work is using the model to evaluate the relative impacts that alternative management strategies can have on overall *B. abortus* transmission.

The National Park Service is exploring the remote delivery of the RB51 brucellosis vaccine to female Yellowstone bison to reduce abortions from this non-native disease and increase tolerance for bison outside YNP (USDI-NPS, 2010). Vaccination is expected to significantly reduce the prevalence of *B. abortus* in bison (Yellowstone Center for Resources,

2008). However, when the parameters of the risk model were modified to evaluate the relative benefits of various management strategies, bison vaccination did not meaningfully reduce *B. abortus* transmission risk to cattle. Management alternatives, such as vaccination, that reduce *B. abortus* prevalence in bison are unlikely to reduce transmission from elk to cattle. These practices would still increase the tolerance for bison outside YNP boundaries, however, as they would decrease either the actual potential for transmission from bison to cattle or the perceived potential for transmission.

Effective strategies included delaying the turn-on date to cattle grazing allotments, reducing elk seroprevalence, reducing the number of cattle at-risk, or prohibiting the comingling of elk and cattle on individual premises. The benefits of a later turn-on date for susceptible cattle provide perhaps the easiest method for reducing the incidence of livestock *B. abortus* infection in the GYA. Combining this strategy with best management practices to reduce comingling on individual premises will reduce the occurrence of cattle brucellosis in the northern GYA. In addition, the reduction in practices that increase elk herd densities and group sizes or the implementation of strategies to reduce elk-to-elk transmission should be promoted to reduce the overall risk to cattle grazing in the northern GYA.

LITERATURE CITED

- RHYAN, J. C., K. AUNE, T. J. ROFFE, D. R. EWALT, S. G. HENNAGER, T. GIDLEWSKI, S. C. OLSEN, AND R. CLARKE. 2009. Pathogenesis and epidemiology of brucellosis in Yellowstone bison: serologic and culture results from adult females and their progeny. *Journal of Wildlife Diseases* 45: 729-739.
- SCHUMAKER, B. A., J. A. K. MAZET, G. E. PLUMB, AND J. VARLEY. 2007. Yellowstone Wildlife Health Program organizational workshop report: June 6-7, 2007. Bozeman, Montana.
- USDI-NPS. 2010. Draft Environmental Impact Statement for Brucellosis Remote Vaccination Program for Bison in Yellowstone National Park. US Department of Interior.
- YELLOWSTONE CENTER FOR RESOURCES. 2008. Vaccination strategies for managing brucellosis in Yellowstone bison. YCR-2008-03, National Park Service, Yellowstone National Park, Wyoming.